

Gluttony and sex in female ixodid ticks: How do they compare to other blood-sucking arthropods?

W. Reuben Kaufman*

Department of Biological Sciences, University of Alberta, Edmonton, Alta., Canada T6G 2E9

Received 24 May 2006; received in revised form 17 September 2006; accepted 2 October 2006

Abstract

The central issue dealt with here is the role of copulation in the control of feeding behaviour in ticks and some haematophagous insects. Female ticks of the family Ixodidae normally engorge to $\sim 100 \times$ their unfed body weight, and then drop from the host, produce and lay eggs, and die. Virgins, on the other hand, normally do not exceed 5–40% (depending on species) of the normal engorged body weight. But instead of detaching voluntarily at that point most virgins remain fixed to the host for extended periods, waiting for males to find them so they can complete engorgement. Virgin haematophagous insects, and virgin ticks of the family Argasidae display little, if any, reduction in blood meal size compared to mated females, at least not during the first ovarian cycle. During subsequent ovarian cycles, meal size in some virgin insects may be somewhat reduced depending on how many eggs are retained in the reproductive tract, but the reduction is not nearly to the same extent as that observed for virgin ixodid females. The stimulatory effect of copulation on engorgement in the latter is caused by a pair of proteins (voraxin α and β) produced in the testis and transferred to the female with the spermatophore. Here, I propose why it might be adaptive for an ixodid female to remain small until mated. The hypothesis is suggested from the facts that ixodid ticks remain attached to the host for days (rather than minutes), and that virgin ticks, above a certain critical weight, lose all opportunity for producing viable offspring should they be groomed off the host prematurely, or should the host die while ticks are still attached.

© 2006 Elsevier Ltd. All rights reserved.

Keywords: Virgin ticks; Ixodidae; Argasidae; Mosquitoes; *Rhodnius*; Tsetse; Bed bugs; Cimex; Feeding behaviour

1. Introduction

Ixodid ticks (Arachnida: Acari: Ixodidae) are obligatory haematophagous arthropods during the three phases of their life cycle—a single larva, a single nymph and adult. Similar to blood-sucking insects, each meal is followed by a moult (larva and nymph) or by spermiogenesis or oogenesis/oviposition. But they differ from blood-sucking insects in at least four important respects.

First, they remain on the host for a number of days (4–14 depending on stage and species) rather than minutes.

Second, although the male takes a relatively small meal (perhaps up to 50% of the unfed body weight), the female is truly gluttonous, increasing her weight approximately 100-fold (Fig. 1). This is at least an order of magnitude

more than the blood-sucking bug *Rhodnius prolixus* and adult ticks of the family Argasidae, about which more will follow.

Third, the final engorged weight actually very much underestimates the true meal size. Adult female *Rhodnius* take the meal (about $3\text{--}4 \times$ the unfed weight) within about 10 min, during which time diuresis begins. The main period of diuresis occurs over the subsequent few hours, eventually resulting in a weight reduction to about twice that of the unfed insect (Maimets, 1985). Although female ixodid ticks likewise excrete the excess fluid of the blood meal, this does not occur during an uninterrupted bout of diuresis following the meal. Instead, interspersed with bouts of blood sucking, the female secretes an enormous volume of hyposmotic saliva back into the host's circulation throughout the feeding period. The blood meal is thus concentrated approximately 2–3-fold while the tick is still attached to the host, and osmoregulation and haemolymph

*Tel.: +1 780 492 1279; fax: +1 780 492 9234.

E-mail address: reuben.kaufman@ualberta.ca.




Fig. 1. Unfed and fully engorged female *Amblyomma variegatum*, demonstrating the astounding size increase (~100-fold). Note that this represents the weight after salivation has disposed of the excess fluid of the blood meal. Hence the total meal volume would be 2–3 times the apparent size of the engorged female in this photo (Koch et al., 1974; Koch and Sauer, 1984). Photo courtesy of Prof. Frans Jongejan, University of Utrecht, The Netherlands, and University of Pretoria, Republic of South Africa, and reproduced with his kind permission.

volume regulation are achieved (Kaufman and Phillips, 1973; Koch et al., 1974; Koch and Sauer, 1984). Hence, the total volume of whole blood removed from the host during a meal is actually somewhere between $200 \times$ and $300 \times$ the unfed female body weight!

Finally, and as will be elaborated further on, the difference in meal size between virgin and mated females is substantially greater than that reported for any insect.

Although the total meal size of ixodid females can be $>200 \times$ their unfed body weight, their feeding progress is not of the slow and steady variety. After *Amblyomma hebraeum* cements itself to the skin and forms the feeding lesion (about a day), it feeds gradually over the next 7–8 days, by which time it will have increased its weight ~ 10 -fold; this is the ‘slow phase’ of engorgement. During the subsequent day (if it has copulated) it increases its weight a further 10-fold (‘rapid phase’ of engorgement). The transition between the slow and rapid phases (at $\sim 10 \times$ the unfed weight) has been called the ‘critical weight’ (CW) because, as reviewed in the next section, important physiological changes occur once the CW is surpassed (Kaufman and Lomas, 1996; Lomas and Kaufman, 1999; Weiss and Kaufman, 2001). With the exception of prostriate ticks (comprising the single genus *Ixodes*), which can copulate before or during the blood meal, copulation in all other ixodid genera (metastriate ticks) occurs only during the feeding period. This is because both sexes among the metastriates require a blood meal to complete gonad maturation (Sonenshine, 1991).

Following engorgement, the salivary glands undergo autolysis within about 4 days, and vitellogenesis proceeds.

Vitellogenin uptake in *A. hebraeum* begins at around 3–4 days post-engorgement, and oviposition begins at around days 10 or 11, and continues for a month or so, after which the spent female dies (Friesen and Kaufman, 2002). An ecdysteroid hormone, probably 20-hydroxyecdysone (20E), triggers both salivary gland autolysis (Harris and Kaufman, 1985; Mao and Kaufman, 1998, 1999) and vitellogenesis (Friesen and Kaufman, 2002, 2004). Injecting high doses of 20E into the haemocoel also inhibits ticks that have been removed from the host below the CW from reattaching (Weiss and Kaufman, 2001). When injected into feeding females, 20E also slows down their feeding rate (Thompson et al., 2005). These observations are consistent with earlier data on ticks that are forcibly removed from the host after exceeding the CW. In such ticks, haemolymph ecdysteroid titre normally rises substantially within a few days, and such large, partially fed ticks generally will not attach to a host if given the opportunity (reviewed by Kaufman and Lomas, 1996 and by Lomas and Kaufman, 1999). It seems likely that this reluctance by large partially fed ticks to reattach and resume feeding is caused by the naturally elevated haemolymph ecdysteroid titre.

2. Mating factors

Seminal fluid is a mixture of secretions from the testis and male accessory glands (MAGs). Gillott (2003) reminds us that the MAG secretions in insects perform numerous functions. MAG secretions are used: (1) to nourish/protect the spermatozoa within the male tract, (2) as components of the spermatophore, (3) to enhance fecundity and attenuate female receptivity, (4) as components of the mating plug, (5) as antimicrobial and antifungal agents to protect the gametes during their sojourn in the female reproductive tract, and (6) as a nutritional supplement to the female. Finally, (7) some MAG secretions extend the duration of sperm viability within the female genital tract. In ticks there are at least four known proteins from the male gonad that are functionally equivalent to MAG proteins (Kaufman, 2004): a sperm capacitation factor (argasid and ixodid ticks) that activates the spermiphores (= spermatozoa) once they are within the seminal receptacle; a vitellogenesis stimulating factor (argasid ticks); a ‘male factor’ (ixodid ticks), that hastens the onset of salivary gland degeneration and ovarian development; and an ‘engorgement factor’ (ixodid ticks) that stimulates the female to engorge. The last two will receive further attention here.

As mentioned above, the salivary glands of female *A. hebraeum* degenerate within about 4 days post-engorgement, a process triggered by 20E. But mated ticks removed from the host at any size above the CW also undergo salivary gland degeneration within 4 days (Harris and Kaufman, 1984), whereas weight-matched virgins above the CW require 8 days (Lomas and Kaufman, 1992a). A protein from the testis, originally named ‘male factor’ by Harris and Kaufman (1984), is the substance responsible

for hastening the rate of salivary gland degeneration from 8 to 4 days, and it achieves this effect by hastening the release of 20E into the haemolymph by 3–4 days (Lomas and Kaufman, 1992b). Vitellogenesis is also triggered by 20E (Friesen and Kaufman, 2002, 2004), so this process is also hastened by ‘male factor’. The engorgement factor will be discussed in the next section.

3. Control of feeding behaviour and the effect of mating on feeding

3.1. Insects

There is a large body of literature relating to the many environmental and physiological factors influencing insect feeding behaviour (Chapman and de Boer, 1995). Not surprisingly, many studies have noted increased feeding rates associated with each ovarian cycle. Females displaying discrete ovarian cycles feed at a high rate during the early part of each cycle, and then feeding rate drops to low levels prior to ovulation (Barton Browne, 1995). Inasmuch as any egg mass produced by virgins is generally smaller than that produced by mated females, it seems inevitable that mated females will feed at a higher rate. Thus, in the milkweed bug (*Oncopeltus fasciatus*) and in the housefly (*Musca domestica*) the oviposition rate of virgins is 30–40% that of mated females, and food intake is less by a comparable amount. Although fewer studies have been conducted on males, a single spermatophore can account for up to 10% of a male’s body weight, and in the male cockroach (*Blattella germanica*), food intake increases as a function of copulation frequency (Barton Browne, 1995).

Until recently, however, it has not been easy to resolve a specific mechanism linking copulation to increased food intake in insects. It is equally surprising that copulation does not seem to enhance feeding in the majority of blood-sucking insects where it has been specifically looked for (see citations in Table 2). What might such discrepancies teach us about the control of feeding behaviour in vector arthropods? There is now one definitive study on how copulation increases feeding rate in a non-vector arthropod: *Drosophila*. Carvalho et al. (2006) demonstrated that mated female *Drosophila* consume over twice as much food within 24 h than do age-matched virgins. Most interesting is that the effect of copulation is mediated specifically by a single MAG peptide (‘sex peptide’ or Acp70A); females mated with males that do not transfer the sex peptide during copulation for one reason or another consume no more than do virgins.

3.2. Ticks

The most outstanding instance of meal size being influenced by mating status is that of female ixodid ticks. As outlined in the Introduction, they normally imbibe $>200\times$ their own unfed weight in host blood over the 7–14-day feeding period, the CW marking the transition

between the slow and rapid phases of engorgement. But this occurs only if the female has copulated. Most virgin *A. hebraeum* do not feed beyond the CW, and can remain attached to the host for weeks waiting for a male to find her. If a male eventually copulates with her, she proceeds to engorge normally; the longer it takes for a male to appear, the more rapidly does the female complete subsequent engorgement (Snow, 1969). Otherwise she remains attached until immunological responses by the host may eventually result in her rejection or death (Brossard and Wikel, 2004). A small percentage of virgin *A. hebraeum* do manage to feed beyond the CW, but only rarely beyond 20% of the engorged weight (Kaufman and Lomas, 1996).

If a tick is removed from the host prematurely, her subsequent behaviour is determined by two circumstances: (a) whether she is above or below the CW, and (b) if above the CW, whether she is virgin or has copulated. Females that are detached from the host below the CW (mated or virgin) maintain a low haemolymph ecdysteroid titre ($<20\text{ ng ml}^{-1}$), do not undergo salivary gland degeneration, or initiate vitellogenesis, and will reattach to a host and continue feeding if given the opportunity. On the other hand, females that are detached from the host above the CW display a rising haemolymph ecdysteroid titre (eventually achieving $300\text{--}500\text{ ng ml}^{-1}$; Kaufman, 1991; Lunke and Kaufman, 1992; Lomas and Kaufman, 1992b), do undergo salivary gland degeneration, do complete vitellogenesis (and ultimately oviposition), and will not reattach to a host and continue feeding if given the opportunity (Kaufman and Lomas, 1996); in all the latter, detached virgins above the CW proceed more slowly than mated females. Finally, in common with some blood-sucking insects, ticks are able to sense an elevated CO_2 concentration in a plume of air—one signal that a potential host is nearby; but detached ticks that have exceeded the CW are no longer responsive to elevated CO_2 (Anderson et al., 1998).

The concept of a CW was established originally over 20 years ago (Harris and Kaufman, 1984). Then, and in subsequent studies, we assessed the CW by any of the several measures just mentioned above. In these studies, we could establish only an approximate value for the CW, because the unfed ticks used, although chosen to be within a fairly narrow weight range, were not individually marked. Weiss and Kaufman (2001), in contrast, marked individual unfed ticks, so that a specific fed-to-unfed weight ratio could be recorded for each individual subsequently removed from the host. In this way we expected to establish a reasonably precise value for the CW. Instead, we discovered that the absolute value of the CW differed depending on which parameter was used to measure it. The CW as defined by reluctance to re-attach to the host was $9\times$ the unfed weight, while that for elevated haemolymph ecdysteroid titre and triggering salivary gland degeneration was $10\times$, for increased ovary weight and oocyte length it was $12\times$, and for increased oocyte vitellin content it was $13\times$ the unfed weight. The biological

meaning of these modest (but significant) differences remains obscure, although it may well relate to a differential sensitivity of each parameter to haemolymph ecdysteroid titre.

What is the mating signal that triggers rapid engorgement in female ixodid ticks? A body of literature, primarily from the 1960s, considered the possibility of mechanical stimuli associated with copulation. However, none of the evidence suggested that a mechanical stimulus (mimicked by inserting inert beads into the seminal receptacle, for example) might be the trigger for rapid engorgement (reviewed by Diehl et al., 1982). Pappas and Oliver (1972) were the first to demonstrate that the male (*Dermacentor variabilis*) produces a specific ‘engorgement factor’ that is transmitted to the female via the spermatophore. (The following sentence contains sexually graphic language; reader discretion is advised!). Females exposed to males whose genital apertures were sealed, did not fully engorge even though they received normal precopulatory stimuli—male mounting behaviour and the male inserting his mouthparts into her gonopore to secrete a droplet of lubricant saliva. Thus, neither mechanical stimulation from the mouthparts, nor a putative signalling molecule from the saliva (e.g.: prostaglandins; Shemesh et al., 1979; Bowman et al., 1996, Aljamali et al., 2002), constitute the engorgement stimulus. In a second experiment, males were exposed to 2.5krad of gamma radiation to kill the spermatids. Females were fed in the company of these irradiated males, and those receiving a spermatophore engorged and oviposited normally; as expected, no embryonic development occurred. Thus, viable spermiophores are not required for full engorgement, but some-

thing else in the spermatophore is Pappas and Oliver (1972).

Weiss et al. (2002) produced 28 recombinant proteins from genes that were upregulated in the testes of fed male *A. hebraeum*. Two of these proteins together were necessary and sufficient for stimulating complete engorgement in virgin females: we named them voraxin α and β (Weiss and Kaufman, 2004). In insects, all of the known male gonad signalling peptides are produced in the MAGs (Wolfner, 2002; Gillott, 2003); voraxin, in contrast, is produced in the testis/vas deferens (Weiss and Kaufman, 2004). Voraxin has not yet been isolated from other tick species, although the following circumstantial evidence suggests that homologous proteins do occur among the Ixodidae: In all species examined to date, virgins feed much less than do mated females, although the unfed-to-fed weight ratio at which virgins stop gaining net weight is variable, ranging from a low of 0.05 for *A. americanum* to a high of 0.35–0.40 for *D. andersoni*, *D. variabilis* and *H. anatolicum* (Table 1, see also Aeschlimann and Grandjean, 1973; Falk-Vairant et al., 1994). Diehl et al. (1982) state that cross-matings between the following species resulted in complete engorgement: *A. americanum* with *A. maculatum*; *Boophilus annulatus* with *B. microplus*; *Rhipicephalus pulchellus* with *R. appendiculatus*; and “...all twelve possible hybrid matings among four species of Kenyan *Amblyomma* ticks (*A. variegatum*, *A. gemma*, *A. eburneum* and *A. cohaerens*).” Whether the recombinant voraxin isolated from *A. hebraeum* is able to stimulate engorgement in other ixodid species remains to be tested, however. Although the concept of a CW (i.e. a weight above which ticks can lay at least some eggs) has been defined in *A. hebraeum*, other

Table 1
Mated and virgin fed weights for a number of ixodid tick species

Species	Engorged weight (mg) mated			Maximum weight (mg) of virgin			Virgin-to-mated weight ratio
	Mean	SEM	N	Mean	SEM	N	
<i>Amblyomma americanum</i>	682	33	9	31.3	6.8	13	0.05
<i>A. hebraeum</i> ^a	2000–2500	—	—	200–250	—	—	0.10
<i>Dermacentor andersoni</i>	791	44	14	267	33	15	0.34
<i>Rhipicephalus sanguineus</i>	212	30	13	36.9	5.4	10	0.17
<i>D. variabilis</i> (from Pappas and Oliver, 1972) ^b	459	—	—	160	27	6	0.35
<i>Hyalomma anatolicum</i> (from Snow, 1969) ^c							
For day 7 ticks	375	—	—	60	—	—	0.16
For day 11 ticks	440	—	—	120	—	—	0.27
For day 14 ticks	415	—	—	160	—	—	0.39

Methods: For each species (*A. americanum*, *D. andersoni*, and *R. sanguineus*), ticks were fed on the shaved backs of rabbits in a divided, foam rubber, cloth-covered arena glued to the back (Kaufman and Phillips, 1973). In one part of the divided arena males were absent (virgin female group), and in the other, an equal number of males was provided (mated female group). Ticks were fed for up to 28 days. All of the mated females engorged (i.e. detached spontaneously) within 12 days (*D. andersoni* and *A. americanum*) or 16 days (*R. sanguineus*). Few virgins ever detached spontaneously; they were removed and weighed after 18 days (*D. andersoni*), after 28 days (*A. americanum*) or after 16 days (*R. sanguineus*). *Ixodes ricinus* and *I. scapularis* were also tested, but most specimens were reluctant to attach to the rabbit and died.

^aRange for *A. hebraeum* is based on extensive data from our laboratory colony. The ratio is based on CW data presented by Weiss and Kaufman (2001).

^bFor Pappas and Oliver (1972) the mean engorged weight (459) was stated without SEM or N.

^cFor Snow (1969), I have estimated the values quoted by inspecting his Fig. 1. In this case, the maximum virgin weight rose linearly with days on the host, whereas the engorged weights on various days appeared not to be significantly different. Thus the virgin-to-mated weight ratio increased steadily.

species have not been examined as carefully. An interesting difference between *Amblyomma* and the other ticks shown in Table 1 is that, in the former, the mean weight at which virgins stop gaining weight is below the threshold for egg-laying. Virgin *Dermacentor* and *Rhipicephalus*, on the other hand, do feed up to a weight at which they lay batches of (infertile) eggs following removal from the host (Pappas and Oliver (1972), and unpublished observations from my laboratory).

4. Comparing ixodid ticks to other vector arthropods

Although the difference in meal size between virgin and mated ixodid female ticks is substantial, this is not the case for other blood feeding arthropods. Potential reasons for this discrepancy will be considered in Section 5.

4.1. Ticks of the family Argasidae

Argasid ticks (vectors for several *Borrelia* species that cause relapsing fever; Sonenshine, 1993) display a lifestyle and reproductive strategy vastly different from that of ixodid ticks. Argasids pass through one larval and up to five nymphal instars, followed by the adult. Males are frequently smaller than females because they emerge from an earlier instar than do females. In *Argas (Persicargas) arboreus*, females usually arise from 3rd or 4th instar nymphs, whereas males usually arise from 2nd instar nymphs (Khalil, 1969). In *Ornithodoros moubata*, males usually emerge from 4th instar nymphs and females from 5th instar nymphs (personal communications from Dr. DeMar Taylor, University of Tsukuba, Japan, and Prof. A. Neitz, University of Pretoria, South Africa). *Ornithodoros*

nymphs and adults feed much more rapidly (30–60 min) than ixodid ticks, though somewhat slower than blood-sucking insects such as *R. prolixus* (~10 min; Maddrell, 1964) or most mosquitoes (as little as 1 min for some anopheline mosquitoes (Chadee and Beier, 1995) or 2–5 min for *Aedes* (Chadee et al., 2002)). The fed weight of adult female *O. moubata* is about 5–12 times their unfed body weight (Diehl et al., 1982). As in insects, excess fluid of the blood meal is excreted to the exterior, although unlike insects, excretion occurs via a pair of coxal organs rather than the Malpighian tubules (Kaufman et al., 1981, 1982). Argasid females can complete a gonotrophic cycle following each meal, laying a modest number of eggs each time. In many of the latter respects, argasid ticks are rather similar to vector insects. Most unlike ixodid ticks (and similar to insects), copulation in *O. moubata*, does not appear to influence meal size much if at all, although it increases rate of digestion and vitellogenesis (Table 2; Aeschlimann and Grandjean, 1973; Germond and Aeschlimann, 1977; Diehl et al., 1982).

4.2. *Rhodnius prolixus*

During the first gonotrophic cycle of this hemipteran, virgin and mated females feed to a similar extent. However, virgin females do lay fewer eggs, thus leaving less room for a subsequent blood meal (Davey, 1997). Consequently, averaged over four cycles (fed every 10 days), virgin females consume 12–22% less blood than mated females (Davey, 1967). Moreover, virgin females use 41–44% more blood to produce a given weight of eggs than do mated females, perhaps because virgins use more of their nutrients in metabolic activity searching for a mate. A far more

Table 2
Average meal sizes in several vector arthropods: a comparison of mated and virgin females

Arthropod	Average meal size (mg) ^a		Virgin to mated ratio	Comments and reference
	Mated females	Virgin females		
Tsetse fly (<i>Glossina palpalis</i>)	29.9	32.1	1.07, NS ^b	Meal 3, females fed every 2 days (Adlington et al., 1996)
Tsetse fly (<i>Glossina palpalis</i>)	34.8	32.6	0.94, NS ^b	Meal 4, females fed every 2 days (Adlington et al., 1996)
Kissing bug (<i>Rhodnius prolixus</i>)	—	—	~1	No difference during first meal, but virgins feed 12–22% less than mated females during subsequent meals (Davey, 1967)
Mosquito (<i>Culex pipiens</i>)	1.94–3.77	1.78–3.17	0.92–0.84, $p < 0.05$	Virgins feed 8–16% less than mated (gravimetric measurement; Adlakha and Pillai, 1976)
Mosquito (<i>Aedes aegypti</i>)	1.44–1.59	1.15–1.29	0.80–0.81, $p < 0.05$	Virgins feed 19–20% less than mated (gravimetric measurement; Adlakha and Pillai, 1976)
Mosquito (<i>Aedes aegypti</i>)	4.3	4.3	1.0	No difference between virgin and mated (spectrophotometric measurement; Klowden, 1979)
Bed bug (<i>Cimex lectularius</i>)	—	—	~1	No difference between virgin and mated (Ussinger, 1966)
Argasid tick (<i>Ornithodoros moubata</i>)	—	—	~1	No difference between virgin and mated (Aeschlimann and Grandjean, 1973; Germond and Aeschlimann, 1977; Diehl et al., 1982).

^aAbsolute meal sizes quoted only when data available for mated and virgin females in the same study.

^bNS: not statistically significant.

complete examination of feeding and mating in *Rhodnius* is presented by Davey in this Special Issue (Davey, 2006).

4.3. *Glossina*

As is the case for many blood-sucking insects, the absolute meal size imbibed by tsetse flies depends on age and on the frequency of feeding (among other factors). In one study using *Glossina palpalis*, Adlington et al. (1996) could detect no significant difference between mated and virgin females fed under the same conditions (Table 2). Males, however, imbibe only about half the amount as virgin or mated females (Loke and Randolph, 1995). These data are from laboratory studies, and probably bear little relevance to what occurs in the field, where copulation normally occurs soon after emergence (Dr. S. Randolph, Oxford University, personal communication). The same may well apply to many insects and argasid ticks that are able to mate prior to taking a blood meal.

4.4. *Mosquitoes*

Adlakha and Pillai (1976) reported that virgin female *Aedes aegypti* imbibe ~20% less blood than do mated females, and that virgin *Culex pipiens fatigans* imbibe 8–16% less blood (Table 2). However, these data were based on gravimetric methods, which are subject to errors because they do not take into account urine losses occurring immediately after the meal. Klowden (1979), measuring haemoglobin content in fed mosquitoes spectrophotometrically (a technique not affected by the extent of diuresis), demonstrated that copulation in *A. aegypti* has no effect on feeding rate (Table 2), although mating apparently does increase the rate of blood meal utilization.

4.5. *Cimex lectularius*

As in *Rhodnius*, adult bed bugs take in relatively smaller meals than the larvae ($1.5 \times$ the unfed weight for the male and $2.2 \times$ for the female; Ussinger, 1966). The blood meal is taken in about 5–10 min, and diuresis begins soon after feeding, half the blood meal volume being excreted within ~5 h. When fed every 12 days, oviposition occurs in a peak/trough cycle, but if fed semi-weekly, egg production becomes essentially continuous, the female putting out more than 5 eggs per week for 4–5 months (Ussinger, 1966). I have found no indication in the literature that mating influences meal size (Table 2); in any case, bed bugs tend to mate after rather than before taking the blood meal (Dr. Ted Morrow, Uppsala University, personal communication).

5. Why does mating have such a profound effect on meal size in female ixodid ticks?

Although the number of vector arthropods referred to in the foregoing is limited, ixodid ticks display significant

differences in lifestyle from all of them. It is tempting to hypothesize that these differences—the necessity for the tick to feed as an adult in order to complete spermiogenesis (with the exception of prostriate ticks), the very long sojourn on the host, the enormous size of the blood meal, and non-nidiculous behaviour (see immediately below)—together may account for ixodid females curtailing full engorgement until mating occurs.

Insects that fly probably encounter a suitable host frequently. Argasid ticks are wingless, but most of them are nidiculous, feeding on the host within the cave, burrow, nest or human-made shelter (Sonenshine, 1993), and so they also have frequent opportunities to feed. In all these cases, it seems to be advantageous to consume numerous but relatively small meals, each triggering a gonotrophic cycle in which relatively few eggs are laid close to home. Because ixodid ticks remain on the host for an extended duration however, one might expect them to drop from the host, remote from home, more frequently. Hence, finding the next meal would be much less certain, and from this one could imagine how imbibing a huge meal, and laying a single, very large batch of eggs (thousands as opposed to tens) might be adaptive. But as outlined just below, the true story is more complicated, because oviposition sites of at least some ixodid ticks can also be close to home.

Ixodid ticks are believed to have evolved from an argasid tick ancestor (Black and Piesman, 1994; Klompen et al., 1996), and some prostriate ticks display nidiculous behaviour (Sonenshine, 1993). The circumstances that might have led the metastriate ixodids to abandon nidiculous behaviour are not known, of course. In this context, however, it is relevant to note that most ixodid ticks tend to complete their engorgement and drop from the host during the scotophase (Sonenshine, 1993), a behaviour that should increase the likelihood for eggs to be laid, and larvae to hatch, near home. A telling exception that supports this rule is the rabbit tick, *Haemaphysalis leporispalustris*, which tend to engorge and detach during the day, when their host is confined to the warren (George, 1971).

Traditionally, the evolutionary pressures on a parasite have been viewed as deriving primarily from the host, but today, other pressures (e.g., habitat/ecological specificity) have gained some currency. For example, many ticks parasitize phylogenetically distinct hosts that share a similar ecological niche. Klompen et al. (1996) reviewed the topic thoroughly and offer the example of *Ornithodoros turicata*, which is found in the burrows of gopher tortoises in the southeastern US. This tick will feed successfully on a number of amphibia, reptiles, birds and mammals that also inhabit these burrows. Some nidiculous ixodid ticks also tend to be distributed near the host's nest. For example, cliff swallows (*Hirundo pyrrhonota*) are known to be parasitized by both the argasid *O. concanensis* and the ixodid *I. baergi* (Sonenshine, 1993). Likewise, *I. uriae* are normally associated with seabird colonies along

North Atlantic coastlines, and all these ticks are most often found in cliff faces and soil near the bird nests (Sonenshine, 1993).

The extended sojourn on the host exposes ixodid ticks to some potential hazards, the two major ones being increased grooming by the host, and activation of the immune system against numerous saliva proteins injected into the host (Brossard and Wikel, 2004; Willadsen, 2004). During the slow phase of engorgement, the tick undergoes significant development and growth in numerous structures—salivary gland (Fawcett et al., 1982), cuticle (Lees and Beament, 1948; Hackman and Filshie, 1982), and reproductive tissues (Diehl et al., 1982)—but relatively modest increase in size ($\sim 10\times$). All this is in preparation for the rapid feeding phase. Remaining small during the slow phase of engorgement might be advantageous because it would seem to be more difficult for a host to dislodge a small tick by grooming. Moreover, because the smaller the tick, the tougher the cuticle, grooming behaviour by the host would also be much less likely to tear open the cuticle and thus kill the tick. These conjectures remain to be formally tested, however.

The final consideration for remaining small as a virgin relates to the facts that parthenogenesis is almost unknown in ixodid ticks (Sonenshine, 1991), and that attaining the CW represents such an important physiological and behavioural milestone. Recall from Section 3 that if ticks are removed from the host prematurely, they are only able to reattach and resume feeding if they are below the CW, and they are only able to lay any eggs at all if they are above the CW (Kaufman and Lomas, 1996; Lomas and Kaufman, 1999; Weiss and Kaufman, 2001). The two main circumstances that would lead to premature detachment in the wild are grooming by the host or death of the host (the latter eventually resulting in the ticks detaching voluntarily). Hence, it is only if the virgin remains below the CW that she would have some chance of finding another host (hopefully infested with some feeding males) and completing the gonotrophic cycle. Once mated, however, ticks removed from the host at any point above the CW are able to lay at least some fertile eggs, the absolute mass being a direct function of the meal size to that point. Such ticks do not have to find another host to complete the gonotrophic cycle. Taken together, it makes sense that virgins should curtail feeding as they approach the CW. If males are not present at that time, they remain attached, but do not feed further. Should they be groomed off the host, or if the host should die, they retain the ability to seek another host. Once males appear and copulate with them, however, they enter the rapid feeding phase (Snow, 1969) and so are now capable of laying at least some fertile eggs if removed prematurely. This hyper-dependence on males for full engorgement has rendered the females of some *Amblyomma* species (*A. hebraeum* is one) reluctant even to attach to an animal that doesn't already harbour feeding males. In our laboratory, for example, we routinely place males on the host at least a day or two before introducing females,

otherwise only a minority of the females will attach within a week.

What accounts for the reluctance of some female *Amblyomma* species to attach in the absence of males? A great deal is now known about the pheromones responsible for aggregation and courtship behaviour in ticks. The reader is referred to Sonenshine (2004) for many further details on the following very brief account. At least four pheromone mixtures lead to courtship in metastriate ixodid ticks, although the details vary considerably among species: (1) An attraction-aggregation-attachment pheromone (AAAP), secreted by some male *Amblyomma* species, stimulates females to congregate and attach near the males. (2) In most other metastriate ticks, feeding females secrete an attractant sex pheromone, 2,6-dichlorophenol, that excites feeding males to detach and search for the attractant source. (3) Feeding females also secrete a non-volatile mounting sex pheromone (cholesteryl oleate in *D. variabilis*; mixtures of different cholesteryl esters in other species) that induces males to probe the females and ultimately leads them to the female's gonopore. (4) A genital sex pheromone (a mixture of long-chain fatty acids and ecdysteroids) is secreted by the vestibular vagina. When the male detects it, he probes the female's genital pore with his mouthparts and ultimately inserts a spermatophore. The genital sex pheromone is the most species-specific of the four, and helps minimize interspecific copulation when several tick species infest the same host.

It is the absence of AAAP that is responsible for some female *Amblyomma* species not attaching to the host unless feeding males are present. (Other metastriate female ticks will attach readily in the absence of males.) AAAP's range of action is the order of a few metres, and its effect is enhanced by carbon dioxide. AAAP is a mixture of primarily three organic volatiles (*o*-nitrophenol, methylsalicylate and non-anoic acid). The concentration ratio among the three substances, and the presence or absence of additional volatile components, vary among species. This variability probably accounts for some species-specific aggregation when multiple species are feeding on the same host, even though each pheromone component may attract all species to some extent. AAAP is secreted by large dermal glands located on the ventral surface of the male (Sonenshine, 2004).

6. Concluding remarks

Ticks do not possess a prothoracic gland, the major source of ecdysone in insects. Instead, in both argasid and ixodid ticks, ecdysone is released from epidermal cells under the control of a neuropeptide (Zhu et al., 1991; Lomas et al., 1997), although it is worth noting that the prothoracic gland is also of ectodermal origin (Klowden, 2002). In some Lepidoptera, the prothoracic gland releases 3-dehydroecdysone, which in turn is converted to ecdysone in the haemolymph by a 3β -reductase enzyme (Warren et al., 1988; Kiriishi et al., 1990). Although 3β -reductase

activity has been detected in the tick salivary gland, it is not found in the haemolymph (Lomas et al., 1998). This is consistent with an earlier observation by Lomas et al. (1997) who did not detect 3-dehydroecdysteroids as secreted products from fragments of integument incubated in vitro; this observation also suggests that the epidermis releases ecdysone directly. Whether ecdysone is released from the epidermis in general, or from a specific population of epidermal cells, has not been determined.

Although voraxin enters the female via the gonopore, we do not know its site of action. The fact that testis homogenates or recombinant voraxin injected into the haemocoel stimulates engorgement in virgin females suggests that voraxin's target is not in the wall of the seminal receptacle. A direct demonstration that voraxin is transported from the seminal receptacle to the haemocoel has to await, however, the development of an ELISA for voraxin. The simplest hypothesis to explain an effect on feeding behaviour is that voraxin stimulates neurons in the synganglion (a single neural mass that comprises the complete CNS in ticks) that innervate the pharyngeal muscles (directly or indirectly). But voraxin and 'male factor' are probably the same proteins, because the two proteins (α and β) constituting voraxin are also necessary and sufficient for hastening salivary gland degeneration and ovarian development in *A. hebraeum* (Weiss and Kaufman, 2004).

An apparent anomaly arises from three additional observations: (1) If partially fed female *A. hebraeum*, below the CW, are forcibly removed from the host, and 20E is injected into the haemocoel before returning them to the host, re-attachment is inhibited for a time; the duration of inhibition is a function of the dose of 20E (Weiss and Kaufman, 2001). (2) Also, when injected into virgin *D. variabilis* ticks while they are attached to the host, exogenous 20E reduces the subsequent rate of blood intake by 21–37% (depending on dose) that of controls (Thompson et al., 2005). (3) Furthermore, the haemolymph ecdysteroid titre of *A. hebraeum* rises significantly during the rapid phase of engorgement (15 ng ml⁻¹ in unfed ticks, 52 ng ml⁻¹ during the rapid phase, falling back to 22 ng ml⁻¹ on the day of engorgement; Mao and Kaufman, 1999). So the highest natural ecdysteroid titre during the attachment period occurs at the very time that feeding rate is most rapid.¹

One might be able to reconcile these apparently contradictory observations as follows. The natural haemolymph ecdysteroid titre during the rapid phase of feeding is still relatively low compared to the exogenous amounts shown to inhibit tick attachment in *A. hebraeum*, and the exogenous amounts shown to inhibit feeding rate in *D. variabilis*. When 20E is injected into the haemolymph, the concentration is initially high, but falls rapidly,

presumably because of two processes: movement from the haemolymph space to the rest of the body (the haemolymph volume of partially fed *A. hebraeum* is approximately 20–25% of body weight; Kaufman et al., 1980) and rapid excretion and/or metabolism (most of the injected ecdysteroid disappears from the haemolymph within 4–7 h; Weiss and Kaufman, 2001). It is thus very difficult to attribute any physiological effect to a unique haemolymph concentration. However, if one uses the initial concentration in the haemolymph following injection, the concentrations at which re-attachment was inhibited were 7.4–384 times the haemolymph concentration that prevails during the rapid phase of engorgement (Weiss and Kaufman, 2001). If one assumes that the haemolymph concentration during the rapid phase of engorgement of *D. variabilis* is similar to that of *A. hebraeum*, the ecdysteroid concentrations at which feeding rate of virgin *D. variabilis* was inhibited were 47–2365 times the haemolymph concentration during the rapid phase of engorgement (calculated from data presented by Thompson et al., 2005). Taken together, the modest rise in ecdysteroid titre occurring naturally during the rapid phase of engorgement would seem to be insufficient to influence feeding rate. The functional significance of the haemolymph ecdysteroid peak during the rapid phase of engorgement, however, is still unknown, although some hypotheses were presented by Lomas et al. (1998). Consequently, voraxin/male factor may have distinct sites of action at the synganglion: one site at which it stimulates engorgement and another at which it hastens the release of the neuropeptide that stimulates ecdysone synthesis in the epidermis. Or indeed, voraxin might act directly on the epidermis to stimulate ecdysone release. Designing experiments to address these questions should not be difficult.

Acknowledgements

Research reported here has been generously funded by an operating grant (Discovery Individual) and a Strategic Projects Grant from NSERC Canada, and an ERATT grant from ASRIP (Alberta Science and Research Investments Program). I am grateful to Ms. Jessica Pollon and Ms. Morenike Olaosebikan for their technical assistance (original data of Table 1). I am most grateful to Dr. Tim Lysyk, Agriculture and Agrifood Canada, Lethbridge Research Centre, Alberta Canada for providing specimens of *D. andersoni*, and Dr. Michael L. Levin, Centres for Disease Control and Prevention, Atlanta, Georgia, USA, for providing specimens of *A. americanum*, *R. sanguineus*, *Ixodes ricinus* and *I. scapularis*. I am most grateful to Prof. Frans Jongejan, University of Utrecht and University of Pretoria, for allowing me to use the image of *A. variegatum* appearing in Fig. 1. Finally, I am grateful to colleagues who offered valuable insights via personal communications: Drs. K.G. Davey (to whom I offer additional thanks for reviewing a complete late draught of the MS),

¹Following engorgement, however, haemolymph ecdysteroid rises much higher, achieving a titre in the range of 300–500 ng ml⁻¹ (Kaufman, 1991; Lunke and Kaufman, 1992; Lomas and Kaufman, 1992b).

C. Gillott, J. Keirans, M. Klowden, T. Morrow, A. Neitz, S. Randolph, S. Simpson, D. Sonenshine and D. Taylor.

References

- Adlakha, V., Pillai, M.K.K., 1976. Role of male accessory gland substance in the regulation of blood intake by mosquitoes. *Journal of Insect Physiology* 22, 1441–1442.
- Adlington, D., Randolph, S.E., Rogers, D.J., 1996. Flying to feed or flying to mate: gender differences in the flight activity of tsetse (*Glossina palpalis*). *Physiological Entomology* 21, 85–92.
- Aeschlimann, A., Grandjean, O., 1973. Influence of natural and “artificial” mating on feeding, digestion, vitellogenesis and oviposition in ticks (Ixodoidea). *Folia Parasitologica (Praha)* 20, 67–74.
- Aljamali, M., Bowman, A.S., Dillwith, J.W., Tucker, J.S., Yates, G.W., Essenberg, R.C., Sauer, J.R., 2002. Identity and synthesis of prostaglandins in the lone star tick, *Amblyomma americanum* (L.), as assayed by radio-immunoassay and gas chromatography/mass spectrometry. *Insect Biochemistry and Molecular Biology* 32, 331–341.
- Anderson, R., Scrimgeour, G., Kaufman, W.R., 1998. Responses of the tick, *Amblyomma hebraeum*, to carbon dioxide. *Experimental and Applied Acarology* 22, 667–681.
- Barton Browne, L., 1995. Ontogenic changes in feeding behaviour. In: Chapman, R.F., de Boer, G. (Eds.), *Regulatory Mechanisms in Insect Feeding*. Chapman & Hall, London, pp. 307–342.
- Black IV, W.C., Piesman, J., 1994. Phylogeny of hard- and soft-tick taxa (Acari: Ixodida) based on mitochondrial 16S rDNA sequences. *Proceedings of the National Academy of Sciences USA* 91, 10034–10038.
- Bowman, A.S., Dillwith, J.W., Sauer, J.R., 1996. Tick salivary prostaglandins: presence, origin and significance. *Parasitology Today* 12, 388–396.
- Brossard, M., Wikel, S.K., 2004. Tick immunobiology. *Parasitology* 129, S161–S176.
- Carvalho, G.B., Kapahi, P., Anderson, D.J., Benzer, S., 2006. Allogenic modulation of appetite by the sex peptide of *Drosophila*. *Current Biology* 16, 692–696.
- Chadee, D.D., Beier, J.C., 1995. Blood-engorgement kinetics of four anopheline mosquitoes from Trinidad, West Indies. *Annals of Tropical Medicine and Parasitology* 89, 55–62.
- Chadee, D.D., Beier, J.C., Mohammed, R.T., 2002. Fast and slow blood-feeding durations of *Aedes aegypti* mosquitoes in Trinidad. *Journal of Vector Ecology* 27, 172–177.
- Chapman, R.F., de Boer, G. (Eds.), 1995. *Regulatory Mechanisms in Insect Feeding*. Chapman & Hall, London.
- Davey, K.G., 1967. Some consequences of copulation in *Rhodnius prolixus*. *Journal of Insect Physiology* 13, 1629–1636 (Hormonal).
- Davey, K.G., 1997. Hormonal controls on reproduction in female Heteroptera. *Archives of Insect Biochemistry and Physiology* 35, 443–453.
- Davey, K.G., 2006. The interaction of feeding and mating in the hormonal control of egg production in *Rhodnius prolixus*. *Journal of Insect Physiology*, this special volume.
- Diehl, P.A., Aeschlimann, A., Obenchain, F.D., 1982. Tick reproduction: oogenesis and oviposition. In: Obenchain, F.D., Galun, R. (Eds.), *Physiology of Ticks*. Pergamon Press, New York, pp. 277–350.
- Falk-Vairant, J., Guerin, P.M., de Bruyne, M., Rohrer, M., 1994. Some observations on mating and fertilization in the cattle tick *Boophilus microplus*. *Medical and Veterinary Entomology* 8, 101–103.
- Fawcett, D.W., Doxsey, S., Buscher, G., 1982. Salivary gland of the tick vector (*R. appendiculatus*) of East Coast fever. II. Cellular basis for fluid secretion in the type III acinus. *Tissue Cell* 13, 231–253.
- Friesen, K.J., Kaufman, W.R., 2002. Quantification of vitellogenesis and its control by 20-hydroxyecdysone in the ixodid tick, *Amblyomma hebraeum*. *Journal of Insect Physiology* 48, 773–782.
- Friesen, K., Kaufman, W.R., 2004. Effects of 20-hydroxyecdysone and other hormones on egg development, and identification of a vitellin-binding protein in the ovary of the tick, *Amblyomma hebraeum*. *Journal of Insect Physiology* 50, 519–529.
- Gillott, C., 2003. Male accessory gland secretions: modulators of female reproductive physiology and behavior. *Annual Review of Entomology* 48, 163–184.
- George, J.E., 1971. Drop-off rhythms of engorged rabbit ticks, *Haemaphysalis leporispalustris* (Packard, 1896) (Acari: Ixodidae). *Journal of Medical Entomology* 8, 461–479.
- Germond, J.-E., Aeschlimann, A., 1977. Influence of copulation on vitellogenesis and egg-laying in *Ornithodoros moubata* Murray (Ixodoidea: Argasidae). In: Adiyodi, K.G., Adiyodi, R.G. (Eds.), *Advances in Invertebrate Reproduction*, vol. 1. Peralam-Kenoth Publishers, Karivellur, Kerala, India, pp. 308–318.
- Hackman, R.H., Filshie, B.K., 1982. The tick cuticle. In: Obenchain, F.D., Galun, R. (Eds.), *The Physiology of Ticks*. Pergamon Press, New York, pp. 1–42.
- Harris, R.A., Kaufman, W.R., 1984. Neural involvement in the control of salivary gland degeneration in the ixodid tick *Amblyomma hebraeum*. *Journal of Experimental Biology* 109, 281–290.
- Harris, R.A., Kaufman, W.R., 1985. Ecdysteroids: possible candidates for the hormone which triggers salivary gland degeneration in the ixodid tick, *Amblyomma hebraeum*. *Experientia* 41, 740–742.
- Kaufman, W.R., 1991. Correlation between haemolymph ecdysteroid titre, salivary gland degeneration and ovarian development in the ixodid tick, *Amblyomma hebraeum* Koch. *Journal of Insect Physiology* 37, 95–99.
- Kaufman, W.R., 2004. Assuring paternity in a promiscuous world: are there lessons for ticks among the insects? *Parasitology* 129, S145–S160.
- Kaufman, W.R., Aeschlimann, A., Diehl, P.-A., 1980. Regulation of body volume by salivation in a tick challenged with fluid loads. *American Journal of Physiology* 238, R102–R112.
- Kaufman, S.E., Kaufman, W.R., Phillips, J.E., 1981. Fluid balance in the argasid tick, *Ornithodoros moubata*, fed on modified blood meals. *Journal of Experimental Biology* 93, 225–242.
- Kaufman, S.E., Kaufman, W.R., Phillips, J.E., 1982. Mechanism and characteristics of coxal fluid excretion in the argasid tick *Ornithodoros moubata*. *Journal of Experimental Biology* 98, 343–352.
- Kaufman, W.R., Lomas, L., 1996. ‘Male Factors’ in ticks: their role in feeding and egg development. *Invertebrate Reproduction and Development* 30, 191–198.
- Kaufman, W.R., Phillips, J.E., 1973. Ion and water balance in the ixodid tick, *Dermacentor andersoni* I. Routes of ion and water excretion. *Journal of Experimental Biology* 58, 523–536.
- Khalil, G.M., 1969. Biochemical and physiological studies of certain ticks (Ixodoidea). Gonad development and gametogenesis in *Argas (Persicargas) arboreus* Kaiser, Hoogstraal and Kohls (Argasidae). *Journal of Parasitology* 55, 1278–1297.
- Kiriishi, S., Rountree, D.B., Sakurai, S., Gilbert, L.I., 1990. Prothoracic gland synthesis of 3-dehydroecdysone and its hemolymph 3 β -reductase mediated conversion to ecdysone in representative insects. *Experientia* 46, 716–722.
- Klompen, J.S.H., Black IV, W.C., Keirans, J.E., Oliver Jr., J.H., 1996. Evolution of ticks. *Annual Review of Entomology* 41, 141–161.
- Klowden, M.J., 1979. Blood intake by *Aedes aegypti* not regulated by insemination. *Journal of Insect Physiology* 25, 349–351.
- Klowden, M.J., 2002. *Physiological Systems in Insects*. Academic Press, New York.
- Koch, H.G., Sauer, J.R., 1984. Quantity of blood ingested by four species of hard ticks (Acari: Ixodidae) fed on domestic dogs. *Annals of the Entomological Society of America* 77, 142–146.
- Koch, H.G., Sauer, J.R., Hair, J.A., 1974. Concentration of the ingested meal in four species of hard ticks. *Annals of the Entomological Society of America* 67, 861–866.
- Lees, A.D., Beament, J.W.L., 1948. An egg-waxing organ in ticks. *Quarterly Journal of Microscopical Science* 89, 291–331.
- Loke, H., Randolph, S.E., 1995. Reciprocal regulation of fat content and flight activity in male tsetse flies (*Glossina palpalis*). *Physiological Entomology* 20, 243–247.

- Lomas, L.O., Gelman, D., Kaufman, W.R., 1998. Ecdysteroid regulation of salivary gland degeneration in the ixodid tick, *Amblyomma hebraeum*: a reconciliation of *in vivo* and *in vitro* observations. *General and Comparative Endocrinology* 109, 200–211.
- Lomas, L.O., Kaufman, W.R., 1992a. The influence of a factor from the male genital tract on salivary gland degeneration in the female ixodid tick, *Amblyomma hebraeum*. *Journal of Insect Physiology* 38, 595–601.
- Lomas, L.O., Kaufman, W.R., 1992b. An indirect mechanism by which a protein from the male gonad hastens salivary gland degeneration in the female ixodid tick, *Amblyomma hebraeum*. *Archives of Insect Biochemistry and Physiology* 21, 169–178.
- Lomas, L.O., Kaufman, W.R., 1999. What is the meaning of ‘critical weight’ to female ixodid ticks?: A ‘grand unification theory’! In: Needham, G.R., Mitchell, R., Horn, D.J., Welbourn, W.C. (Eds.), *Acarology IX*, vol. 2, *Symposia*. The Ohio Biological Survey, Columbus, OH, pp. 481–485.
- Lomas, L.O., Turner, P.C., Rees, H.H., 1997. A novel neuropeptide-endocrine interaction controlling ecdysteroid production in ixodid ticks. *Proceedings of the Royal Society of London B* 264, 589–596.
- Lunke, M.D., Kaufman, W.R., 1992. Effects of the avermectin analogue MK-243 on vitellogenesis and reproduction in the ixodid tick, *Amblyomma hebraeum*. *Experimental and Applied Acarology* 13, 249–259.
- Maddrell, S.H.P., 1964. Excretion in the blood-sucking bug, *Rhodnius prolixus* Stal. II. The normal course of diuresis and the effect of temperature. *Journal of Experimental Biology* 41, 163–176.
- Mao, H., Kaufman, W.R., 1998. DNA binding properties of the ecdysteroid receptor in the salivary gland of the female ixodid tick, *Amblyomma hebraeum*. *Insect Biochemistry and Molecular Biology* 28, 947–957.
- Mao, H., Kaufman, W.R., 1999. Profile of the ecdysteroid hormone and its receptor in the salivary gland of the adult female tick, *Amblyomma hebraeum*. *Insect Biochemistry and Molecular Biology* 29, 33–42.
- Maimets, Ilo, 1985. The transport of a blood meal through the alimentary canal of *Rhodnius prolixus*. M.Sc. Thesis, York University, Toronto, Canada, 95pp.
- Pappas, P.J., Oliver Jr., J.H., 1972. Reproduction in ticks (Acari: Ixodoidea). 2. Analysis of the stimulus for rapid and complete feeding of female *Dermacentor variabilis* (Say). *Journal of Medical Entomology* 9, 47–50.
- Shemesh, M., Hadani, A., Shklar, A., Shore, L.S., Meleguir, F., 1979. Prostaglandins in the salivary glands and reproductive organs of *Hyalomma anatolicum excavatum* Koch (Acari: Ixodidae). *Bulletin of Entomological Research* 69, 381–385.
- Snow, K.R., 1969. Egg maturation in *Hyalomma anatolicum anatolicum* Koch 1844 (Ixodoidea, Ixodidae). In: Owen Evans, G. (Ed.), *Proceedings of the Second International Congress Acarology*. Akademiai Kiado, Budapest, pp. 349–356.
- Sonenshine, D., 1991. *The Biology of Ticks*, vol. 1. Oxford University Press, Oxford.
- Sonenshine, D., 1993. *The Biology of Ticks*, vol. 2. Oxford University Press, Oxford.
- Sonenshine, D., 2004. Pheromones and other semiochemicals of ticks and their use in tick control. *Parasitology* 129, S405–S425.
- Thompson, D.M., Khalil, S.M.S., Jeffers, L.A., Ananthapadmanaban, U., Sonenshine, D.E., Mitchell, R.D., Osgood, C.J., Apperson, C.S., Roe, R.M., 2005. *In vivo* role of 20-hydroxyecdysone in the regulation of the vitellogenin mRNA and egg development in the American dog tick, *Dermacentor variabilis* (Say). *Journal of Insect Physiology* 51, 1105–1116.
- Ussinger, R.L., 1966. *Monograph of Cimicidae*. The Thomas Say Foundation, vol. VII. Entomological Society of America.
- Warren, J.T., Sakurai, S., Rountree, K.B., Gilbert, L.I., Lee, S.S., Hakanishi, K., 1988. Regulation of the ecdysteroid titer of *Manduca sexta*: reappraisal of the role of the prothoracic glands. *Proceedings of the National Academy of Sciences, USA* 85, 958–962.
- Weiss, B., Kaufman, W.R., 2001. The relationship between ‘critical weight’ and 20-hydroxyecdysone in the female ixodid tick, *Amblyomma hebraeum*. *Journal of Insect Physiology* 47, 1261–1267.
- Weiss, B.L., Kaufman, W.R., 2004. Two feeding-induced proteins from the male gonad trigger engorgement of the female tick, *Amblyomma hebraeum*. *Proceedings of the National Academy of Sciences USA* 101, 5874–5879.
- Weiss, B.L., Stepczynski, J.M., Wong, P., Kaufman, W.R., 2002. Identification and characterization of genes differentially expressed in the testis/vas deferens of the fed male tick, *Amblyomma hebraeum*. *Insect Biochemistry and Molecular Biology* 32, 785–793.
- Willadsen, P., 2004. Anti-tick vaccines. *Parasitology* 129, S367–S387.
- Wolfner, M.F., 2002. The gifts that keep on giving: Physiological functions and evolutionary dynamics of male seminal proteins in *Drosophila*. *Heredity* 88, 85–93.
- Zhu, X.X., Oliver Jr., J.H., Dotson, E.M., 1991. Epidermis as the source of ecdysone in an argasid tick. *Proceedings of the National Academy of Sciences of the USA* 88, 3744–3747.