



SMITHSONIAN INSTITUTION
U. S. NATIONAL MUSEUM

Vol. 86

Washington : 1939

No. 3048

REVISION OF THE BEETLES OF THE MELOLONTHINE
SUBGENUS PHYTALUS OF THE UNITED STATES

By LAWRENCE W. SAYLOR

SEVERAL years ago I began the study of the scarabæid beetles of the subgenus *Phytalus* Erichson (genus *Phyllophaga* Harris) with the aim of bringing together in one paper data on all the described species, since the literature on the group is somewhat scattered and unavailable to many. It develops that several changes in taxonomic standing are necessary, and the knowledge of the distribution of the various species is greater than has been recorded. In this subgenus the genitalia of both sexes are of great use as supplementary diagnostic characters, and except in the case of *Phyllophaga* (*Phytalus*) *amani* Sanderson they have not been previously figured.

I am indebted to Dr. E. A. Chapin, Dr. M. A. Sanderson, Prof. E. C. Van Dyke, and Mark Robinson for the loan of material and for many other kindnesses.

The group as now defined is restricted to the American continents and adjacent islands and includes more than 60 described species, more than half of which inhabit Central America and the West Indies. In the United States the insects are found most commonly in the southern regions, specimens having been seen from Arizona, New Mexico, Texas, Georgia, Alabama, and Tennessee and also from New Jersey. I have recently received specimens of two of our United States species, *P. pallida* Horn and *P. sonora* Saylor (= *debilis* LaConte), from northern Mexico; a check of the literature reveals that these two species are apparently not recorded under any other name in Mexican faunal works.

107283—39

157

Saylor, Lawrence W. 1939. Revision of the beetles of the Melolonthine subgenus *Phytalus* of the United States. 86:3048. 157-167.

Phytalus vexatus Horn (= *cavifrons* Linell) has been removed by Chapin to the genus *Chlaenobia*, which differs from *Phyllophaga* (and *Phytalus*) in that the tarsal soles are usually very densely pilose in the male and the hind thoracic angles usually constricted rather abruptly at the base; the genus *Chlaenobia* also has a distinct facies, enabling one familiar with the group to recognize specimens on sight. Some of the true *Listrocheilus* (e. g., *scoparius*) also have densely pilose soles in the male sex, and later studies may show that *Chlaenobia* may be better placed as a subgenus.

Phytalus is no longer tenable as a genus because some of the species exhibit characters that absolutely grade into those of *Phyllophaga* and leave no single character or group of characters distinguishing the two. Although it can no longer be used in a generic sense, the name *Phytalus* is of use subgenerically as applied to a group of species having narrowly cleft claws in at least the male sex.

In probably no other group of American scarab beetles is there such a variety of opinion as to the validity of genera and of their names as in that of the phyllophagans. In 1920 Arrow, of the British Museum, stated that *Phytalus*, *Brahmina*, and *Holotrichia* were not separable from *Phyllophaga* (= *Lachnosterna*), and with this I heartily agree,¹ as I have found by experience that the use of the claws alone for generic characters is in most instances unsatisfactory because the claws may be very different in species that in every other character are obviously of the same genus.

In *Phyllophaga*, as at present recognized, there are several good groups, which, if segregated as different genera, however, would separate species greatly alike in most characters and apparently of the same lineage. Such a separation would, in most instances, be on the basis of the male characters alone, certainly undesirable criteria for generic definitions. Among such characters are the deformed middle claws, fixed hind tibial spurs, and narrowly cleft tarsal claws; the first two are nonvariable, but the last varies greatly in degree in the sexes. Another group could be defined if the very long basal claw dilation, giving the appearance of a third tooth, were used; in the single species concerned, *P. heteronycha* Bates, the tarsal claws of the fore and midlegs are 3-toothed, while those of the hind pair, like those of *Phytalus*, are narrowly cleft. Such a segregation is unadvisable, however, as all degrees in length of the tooth formed by the basal dilation can be found in various species. Furthermore, in my opinion it would serve no useful purpose to form a special genus for those species in which the usual three segments of the antennal club are increased to four or five, as other characters are the same as in the species with the normal number of segments in the club.

In the course of the present studies, I examined nearly every one of the United States species and also well over a 100 species of Neotropical *Phyllophaga* and allied genera, and in addition more than 50 species of Oriental *Brahmina*, *Holotrichia*, and allied groups. The necessity is apparent of suppressing the name *Brahmina* entirely and of ranking the American *Phytalus* and the Oriental *Holotrichia* as subgenera of *Phyllophaga*. This action has been suggested at one time or another, in whole or in part, by nearly every serious student of the group from Blanchard's time on (Blanchard, Bates, Arrow, Chapin, and Saylor), and the change was finally made by me in a recent paper (see footnote 1).

The position of other related genera, *Listrochelus* and *Chirodines*, is also open to some question; the former name seems to be valid for subgeneric use if restricted to a certain group of species, as has been done in a revision now in preparation by Dr. E. A. Chapin and myself. The genus *Chirodines* was separated on the basis of only a slight difference in the claws, and when thoroughly studied may quite possibly be shown to merit only subgeneric status, or may entirely fail of recognition.

Little is known regarding the economic status of the majority of the species, but *P. pallida* Horn has been observed doing a good deal of damage in Arizona by eating the foliage of rose bushes, young fruit trees, and walnut trees, often stripping them.

If the subgenus *Phytalus* is restricted to those species of *Phyllophaga* having the very narrowly cleft tarsal claws, it embraces the following species in our fauna: *P. bilobatata* Saylor, *P. georgiana* Horn, *P. omani* Sanderson, *P. pallida* Horn, *P. sandersonia* Saylor, *P. sonora* Saylor, and *P. obsoleta vanalleri* Schaeffer. The males usually have a flat or convex abdomen, as viewed from the side, and the antennal club is as long or nearly as long as the funicle; the females usually have the abdomen concave and robust and the antennal club much shorter than the funicle.

KEY TO THE SPECIES OF THE SUBGENUS PHYTALUS

MALES

1. Hind tibiae with 1 spur free, the other fixed and immovable..... 2
Both hind tibial spurs free, movable..... 4
2. Upper tooth of claw longer to very much longer than lower
one; antennae 9-segmented..... 3
Upper tooth much shorter and narrower than lower tooth;
antennae 10-segmented..... sandersonia
3. Yellow-testaceous; scutellum distinctly punctured only at sides;
lower tooth of claw two-thirds or more as long as upper;
fixed spur of hind tibia directly contiguous with tibial margin..... omani
Rufotestaceous; scutellum evenly punctured over entire surface;
lower claw tooth shorter; not more than one-half as long as
upper; entire apical margin or posterior tibia distinct..... georgiana
4. Fifth abdominal segment flattened at middle and coarsely
punctate, not granulate; color reddish brown to piceous..... bilobatata
Fifth abdominal segment lobate or with granular tumosities..... 5
5. Upper tooth of claw much shorter than lower, claw very wide
at base; fifth abdominal segment with a large triangularly
shaped lobe, the latter incised at apex and projecting back
over sixth segment..... obsoleta vanalleri
Upper tooth of claw much longer than lower, fifth abdominal
segment without a triangular lobe..... 6
6. Lobe of fifth abdominal segment reaching to or beyond apical
margin, granulate, faintly bilobed at apex; pygidium coarsely
rugose-punctate..... pallida
Lobe of fifth abdominal segment less evident, the surface more
tumid, with transverse granules; pygidium smooth, sparsely
punctured..... sonora

FEMALES

1. Pygidium with a distinct tubercle just above apex..... 2
Pygidium without tubercle, plane or sometimes thickened api-
cally..... 3
2. Densely clothed above with short, erect, tawny hairs; thorax
regularly and densely punctured, the punctures practically
touching one another..... sandersonia
Pronotum and elytra nearly or quite glabrous; thorax sparsely
punctured, the punctures separated by 2 to 4 times their di-
ameters..... obsoleta vanalleri
3. Lateral thoracic margins distinctly crenulate; claws narrowly
cleft, upper tooth longer than lower; front very closely,
coarsely rugose-punctate..... pallida
Lateral thoracic margins entire or nearly so..... 4
4. Clypeus narrowly and deeply emarginate; color dark castaneous
to rufopiceous..... bilobatata
Clypeus broadly but not deeply emarginate; color testaceous or
rufotestaceous..... 5
5. Last abdominal segment almost flat, slightly transversely sul-
cate; pygidium densely or sparsely punctured (Eastern
United States)..... 6

Last abdominal segment convex, not sulcate; pygidium sparsely punctured (Arizona, Mexico)-----	sonora
6. Scutellum densely, closely punctured; pygidium evenly punctured-----	georgiana
Scutellum punctured only at sides; pygidium irregularly punctured-----	omani

PHYLLOPHAGA (PHYTALUS) SANDERSONIA, new name

PLATE 9, FIGURES 1a-1c

Phytalus robustus HORN, Trans. Amer. Ent. Soc., vol. 12, p. 120, 1895 (*nec* LeConte, 1856).

Phytalus trichodes BATES, Biologia Centrali-Americana, vol. 2, pt. 2, p. 178, 1890.

Male.—Robust, elongate-oval, rufocastaneous, shining, entirely clothed above with short, very dense, suberect hair, that of front longer. Head with front exceedingly densely variolate-punctate, the punctures of moderate size and closely contiguous; clypeus punctured like front but less densely so, its apex moderately reflexed and slightly emarginate at middle. Antennae 10-segmented, club subequal to or very slightly longer than funicle. Thorax very regularly and extremely densely, evenly punctured over the entire surface; hind angles obtusely angulate; lateral margins almost evenly arcuate, crenulate, ciliate. Elytra very rugosely wrinkled, densely punctured, striae other than sutural weakly indicated. Pygidium with or without a very short longitudinal carina at center of basal margin, surface flattened near apex, evenly and densely punctured over the entire surface, with dense moderately long suberect hairs; apex subtruncate to subrounded. Abdomen polished, sparsely hairy, widely, shallowly and longitudinally concave at center; fifth segment plane, with moderately dense setigerous punctures at sides and apex; sixth two-thirds as long as fifth, more coarsely punctured and with longer erect hairs. Fixed spur of hind tibia short and twisted. All claws with upper tooth much shorter and more slender than lower. Front tarsi with segments 1 to 3, inclusive, with the inner apical margin prolonged into a strong spine, this character most strongly marked in segment 1.

Female.—Pygidium glabrous, sparsely punctured, declivate, and with a tubercle before apex; posterior tibial spurs free, elongate. (Amended from original description.)

Length, 17-20 mm. Width, 8.5-10 mm.

Remarks.—Horn described the species from the Rio Grande country near Matamoros, and I have seen two specimens (that were compared with the type) from Brownsville, Tex. (F. H. Snow and Charles Schaeffer), and also one male from Del Rio, Tex. (May 15, 1937, A. Meade). Described by Bates from Las Vigas, Veracruz

Mexico, and seen by me from Jalapa and Monterrey in Mexico (June). A comparison of the external features and of the genitalia of a cotype of Bates' species with Horn's species proves the two are identical. This species, to date, has been very rare in the United States, and less than a dozen specimens in American collections are known to me as having been taken within our boundaries; it was cited by Bates as being rather numerous at Las Vigas, Veracruz. The species is named for my good friend Dr. M. W. Sanderson.

PHYLLOPHAGA (PHYTALUS) OMANI Sanderson

PLATE 9, FIGURES 2a-2e

Phyllophaga (Phytalus) omani SANDERSON, Journ. Kansas Ent. Soc., vol. 10, p. 66, 1937.

Male.—Elongate, cylindrical, yellow-testaceous, the head piceous, surface shining. Head with front moderately and not closely punctured, the punctures separated by one to several times their own diameters; clypeus deeply marginate and moderately punctured; clypeal suture lightly impressed. Antennae 9-segmented, the club as long as the entire stem, unicolorous. Prothorax moderately, rather unevenly punctured, the punctures closer along the anterior and basal margins; sides parallel in basal half, then gradually rounded to apex; with an indistinct fuscous spot near the rounded lateral margins. Elytra punctured like thorax, somewhat rugose, costae except sutural obsolete. Pygidium rather strongly convex, finely and evenly punctured with a few short hairs at apex, remaining surface glabrous. Abdomen somewhat flattened at middle and with vague longitudinal impressions on segments 3 and 4; segment 5 plane; segment 6 slightly excavated at middle and with a small punctate elevation each side of middle. One of the spurs of the hind tibia short and fixed. Upper portion of the claw nearly as wide as lower and distinctly longer. Lower claw margin very finely and minutely, irregularly crenulate.


Female.—Club of antennae shorter than funicle; hind tibial spurs free; abdomen somewhat flattened at middle; pygidium transverse and evenly punctured. Otherwise similar to male.

Length, 14 mm. Width, 6 mm.


EXPLANATION OF PLATES 9 AND 10

1, *Phyllophaga (Phytalus) sandersonia*, new name; 2, *P. (P.) omani* Sanderson; 3, *P. (P.) georgiana* (Horn); 4, *P. (P.) bilobatata*, new name; 5, *P. (P.) pallida* (Horn); 6, *P. (P.) sonora*, new name; 7, *P. (P.) obsoleta vanalleri* (Schaeffer).


Letters indicate views, as follows: *a*, En face view of male genitalia; *b*, side view of male genitalia; *c*, tarsal claw; *d*, female genitalia; *e*, en face-ventral view of male genitalia.




7 a




7 b




7 c




4 a




4 c



4 b




6 a




6 b




6 c



3 a



3 b




3 c



5 a




5 c



5 b

GENITALIA AND TARSAL CLAW OF PHYTALUS.

(FOR EXPLANATION OF PLATE SEE PAGE 162.)



GENITALIA AND TARSAL CLAW OF PHYTALUS.

(FOR EXPLANATION OF PLATE SEE PAGE 162.)

Remarks.—This interesting species is known only from the type material; the above description is condensed from the original. The holotype and allotype are from Burnsville, Ala. (July 20, 1930), and Prattsburg, Ga. (July 24, 1930), respectively; through the courtesy of Dr. Sanderson I was allowed to examine the male type before its description. One male in my collection from "North America."

PHYTALOPHAGA (PHYTALUS) GEORGIANA (Horn)

PLATE 9, FIGURE 3*d*; PLATE 10, FIGURES 3*a*–3*c*

Phytalus georgianus HORN, TRANS. AMER. ENT. SOC., vol. 12, p. 122, 1885.

Male.—Rufous to rufotestaceous, head and thorax darker, surface shining with a few short scattered hairs. Front of head coarsely punctured, slightly rugose, with a small very irregular impunctate area near the vertex; clypeal suture deeply impressed at sides only; clypeus coarsely rugose-punctate; apex acutely not deeply but rather broadly emarginate; antennal club slightly longer than remainder of the antennae, third and fourth segments elongate. Sides of thorax not crenulate, hind angles rectangular, front angles very obtuse, disk coarsely and densely variolate-punctate, a few minute hairs near middle of the base. Elytra densely and coarsely rugose-punctate. Pygidium very convex, rather densely and irregularly punctured, glabrous, apex broadly rounded. Abdomen flattened at middle, polished, with a few setigerous punctures; segment 5 much shorter than 6, abruptly and narrowly declivate at apical margin; segment 6 slightly concave at center, with a transverse carina, the latter bearing a row of sparse hairs, and the row interrupted at middle. Hind tibial spurs spiniform, with the fixed one half as long as the other.

Female.—In the specimen at hand, the antennae are 8-segmented, with segment 3 very elongate, but it is probable that the normal number of segments is 9; club equal to segments 3–5 combined; last abdominal segment very shallowly transversely sulcate, sparsely punctured and fimbriate at apex; tibial spurs elongate, free; otherwise similar to male.

Length, 12.5–13 mm. Width, 6–6.5 mm.

Remarks.—I have seen collected specimens from Whitesbog, N. J. (July 9), "Barcoure," Ala., and three bred specimens from Lakehurst, N. J.

Described from Georgia, this rather rare species is distinctly separated from the others by the sexual characters. Horn, in describing the species, apparently overlooked the fact that one hind tibial spur in the male is definitely fixed and immovable, though unless careful examination is made the spurs appear to be free.

PHYLLOPHAGA (PHYTALUS) OBSOLETA VANALLERI (Schaeffer)

PLATE 9, FIGURE 7*d*; PLATE 10, FIGURE 7*a-7c*

Phytalus obsoletus BLANCHARD, Catalogue de la collection entomologique, vol. 1, p. 131, 1850.

Phytalus vanalleri SCHAEFFER, Bull. Brooklyn Ent. Soc., vol. 22, p. 215, 1927.

Male.—Elongate, subparallel, head and thorax rufous, otherwise testaceous to rufotestaceous, thorax and elytra glabrous, shining. Front of head closely, more or less confluent punctured, with long, erect, testaceous hair; a narrow area behind clypeal suture smooth and impunctate; clypeal suture not impressed; clypeus broad, faintly emarginate at middle of the subtruncate apex, the latter strongly reflexed, disk coarsely and sparsely punctured; antennal club usually longer than the rest of the antennae. Thoracic angles obtuse but well defined, base margined except at the middle, sides subangulate, subcrenulate; disk moderately, densely, and regularly umbilicate-punctate, with a small, irregular, median impunctate area; a few of the punctures with a minute testaceous hair. Elytra densely rugose-punctate, with a few hairs near apex. Pygidium very convex, polished, sparsely punctured, with suberect hairs; disk narrowly impressed before apex, the latter broadly rounded and ciliate. Abdomen shallowly impressed at middle, the concavity densely and finely setigerously punctate, the hair long and fine; segment 5 with a broad transverse carina near apical margin widely interrupted at middle, the carina densely pilose; segment 5 including the lobe as long as segments 3 and 4 combined; sides of abdomen sparsely punctured. Posterior spurs long, free.

Female.—Median impunctate area of thorax broader, with the punctures more distinct, middle of abdomen very slightly canalicate, regularly and sparsely punctured, almost glabrous and highly polished; segment 5 more densely punctured at the slightly raised apical margin; segment 6 slightly convex, coarsely and sparsely punctured; antennal club equal to segments 3-7 combined; otherwise similar to male.

Length, 16-17 mm. Width, 7.5-8 mm.

Remarks.—This form is rather uncommon in the Southeast, but specimens have been seen from Alabama, Texas, and Louisiana. It is easily separated from all others of our fauna by the sexual characters, and in these it is similar to *P. obsoleta* Blanchard, which is a common species in Mexico, Guatemala, and Nicaragua; the two are very close, and *P. vanalleri* Schaeffer is probably best regarded at present as a northern subspecies having a longer antennal club (often two-fifths longer than the funicle), more crenate lateral thoracic margins, less elevated pectinate teeth on the lobe of the fifth abdominal

segment, very sparsely punctured clypeus, and smoother front; individuals are not lacking, however, in which almost all intermediate degrees of these characters appear, and possibly when more specimens have been seen from northern Mexico and southeastern United States the name *P. vanalleri* Schaeffer may have to be withdrawn altogether.

The antennal club of *P. vanalleri* Schaeffer is very variable; in most specimens it is as long as or longer than the stem and scape combined, while in almost all specimens of typical *P. obsoleta* Blanchard the club is noticeably shorter than the scape and funicle combined (averages about one-fifth longer than the funicle) as well as being lighter in color. The clypeus of typical *P. obsoleta* Blanchard is almost always densely punctured, while that of *P. vanalleri* Schaeffer is sparsely punctured at the middle of the clypeal suture.

PHYLLOPHAGA (PHYTALUS) BILOBATATA, new name

PLATE 9, FIGURE 4d; PLATE 10, FIGURES 4a-4c

Phytalus cephalicus HORN, Trans. Amer. Ent. Soc., vol. 12, p. 120, 1885 (not *cephalica* LeConte, 1856).

Male.—Elongate, dark castaneous to piceous, shining, usually glabrous above. Head with front densely variolate-punctate; clypeal suture rather deeply impressed, slightly bisinuate; clypeus broad, deeply and very narrowly emarginate at apex, giving a bilobed appearance to the apical margin, surface slightly tumid in some examples, densely, coarsely, and somewhat confluent punctured; antennal club slightly smaller than the stem, antennae 10-segmented. Thorax with a faint suggestion of a longitudinal sulcus on the disk in some examples, sides obtusely rounded, hind angles obtuse but distinct, margin entire, disk finely, sparsely and rather regularly punctured, a more or less irregular smooth space at middle. Elytra moderately densely and rugosely punctured, sometimes with very minute testaceous hairs. Pygidium very convex, densely and rugosely punctured, the punctures each with a short hair; apex broadly rounded, with a few longer hairs. Abdomen convex, very sparsely and finely punctured at middle, segment 5 depressed behind, a small group of sparse, long, erect hairs on each side of the segment; segment 6 elevated, densely punctate, abruptly declivous at base, in some cases with a faint trace of a longitudinal impression. Posterior spurs long, narrow, and free.

Female.—Antennal club much smaller; abdomen more convex, last segment longer and less densely punctured; otherwise similar to male.

Length, 15-18 mm. Width, 6-7.5 mm.

Remarks.—All positively identified material is from Arizona; localities represented are Nogales, Carr Canyon, Fort Grant, and "Chiri-

cahua Mts.," all collected in June or July. The pygidium and clypeus vary from slightly convex to tumid; in one female example the basal half of the pygidium is semitumid. Schaeffer points out that the last paragraph in Horn's description of *P. robusta* (Horn) (i. e., *sander-sonia* Saylor) in reality refers to this species, which would seem to indicate that this species occurs in New Mexico; I have not seen specimens, however, from that State.

PHYLLOPHAGA (PHYTALUS) PALLIDA (Horn)

PLATE 9, FIGURE 5*d*; PLATE 10, FIGURES 5*a*-5*c*

Phytalus pallidus HORN, Trans. Amer. Ent. Soc., vol. 12, p. 121, 1885; 6th Ann. Rept. Arizona Comm. Hort. and Ent., p. 30, 1914.

Male.—Testaceous, head and thorax rufotestaceous, almost glabrous, surface moderately shining. Head with front densely and coarsely punctured; clypeal suture lightly impressed, bisinuate; clypeus slightly emarginate at center of apex, faintly reflexed, angles rounded, surface densely punctured; antennal club slightly shorter than funicle. Thorax with sides broadly rounded, crenulate, angles obtuse, disk coarsely sparsely and rather regularly punctured. Elytra densely punctured, rugose, sutural costae elevated. Pygidium flattened, coarsely rugose, moderately and densely punctured, glabrous except for a few erect hairs at the rounded apex. Abdomen finely and moderately densely punctured, the punctures with short hairs; apical half of segment 5 with a raised rounded lobe reaching to apical border, the surface of which is granular-strigose, its apex finely serrate; segment 6 rather deeply and transversely sulcate, the apical and basal margins carinate. Posterior spurs free, elongate.

Female.—Antennal club equal to segments 3-7 combined; abdomen convex, sparsely setigerously punctate, segment 5 longer than 4, slightly tumescent in apical half, densely and coarsely punctured; segment 6 one-half the length of 5, densely punctured; otherwise similar to male.

Length, 12-14 mm. Width, 5.5-6 mm.

Remarks.—Most of the specimens examined are from Arizona: Fort Huachuca, Fort Grant, and Ramsey Canyon, all taken in July. I have also in my collection a male from Bakachaka, Rio Mayo, Sonora, Mexico, taken in July by my friend Howard Gentry.

Closely related to *P. sonora* Saylor but may be separated by the puncturing of the head and clypeus, as well as by the male sexual characters.

PHYLLOPHAGA (PHYTALUS) SONORA, new name

PLATE 9, FIGURE 6*d*; PLATE 10, FIGURES 6*a*–6*c**Phytalus debilis* HORN, Trans Amer. Ent. Soc., vol. 12, p. 122, 1885 (not LeConte, 1856).

Male.—Highly polished, glabrous, rufotestaceous, varying at times almost to piceous. Head with front coarsely and moderately densely punctured, the punctures shallow, more or less umbilicate, usually separated by one to two times their diameters, a large impunctate area on vertex; clypeal suture impressed, strongly biarcuate; clypeus wide, relatively flat, apex narrowly emarginate, slightly reflexed, disk coarsely and sparsely punctured. Antennal club longer than funicle. Thorax with sides entire, obtusely rounded at middle, slightly sinuate behind, front and hind angles obtuse, the latter prominent; disk regularly punctured, the punctures separated by one to three times their diameters. Elytra sparsely and shallowly punctured. Pygidium flattened, sides slightly concave, very sparsely punctured, glabrous except for a few erect hairs on margin near the subtruncate apex. Abdomen slightly flattened and very sparsely punctured at middle; posterior half of segment 5 at center with a raised granulate lobe, the latter not quite reaching the apical margin of the segment, segment 6 narrowly and transversely impressed, with a row of cilia along apical margin. Posterior spurs free.

Female.—More robust, clypeal suture more deeply impressed, faintly biarcuate; pygidium just before apex with a smooth slightly raised area, the apex of which has a single row of rather large punctures, each with a long erect hair; spurs long, apices rounded; abdomen convex, with segment 5 finely punctured in apical half, segment 6 somewhat transversely impressed at base, sparsely punctured, apex ciliate; the 10-segmented unicolorous, dark antennae with the club elongate, equal to segment 3–7 combined; hind femora quite broad; otherwise similar to male.

Length, 10–14 mm. Width, 5.5–6 mm.

Remarks.—Most of the material is from Arizona: Tucson, Carr Canyon, Globe, Patagonia, and Badger, taken in June, July, and August. I have a dozen examples in my collection from Mexico, all collected in Rio Mayo, Sonora, by Howard Gentry, at Sierra Charuca (July), San Bernardo (July), and Vinaterio (June). The species has not been recorded from Mexico before.

A not uncommon species of which the female has apparently not previously been recognized. In my collection is one individual of that sex from San Bernardo, Mexico (Gentry), from which the diagnosis has been drawn. The male pygidium may vary from almost flat to rather strongly convex. In some few examples, the sides of the thorax are very finely crenulate.