

CATALOGO DE LOS DIPTERA
DE NICARAGUA.
4. CULICIDAE (NEMATOCERA).

Por Jean-Michel Maes *
& Pedro Rivera Mendoza.**

Resumen.

Este catálogo presenta las 40 especies de Culicidae (Diptera : Nematocera) reportadas de Nicaragua. Para cada especie se cita la sinonimia, la distribución geográfica, los hospederos, las enfermedades transmitidas y los enemigos naturales. La bibliografía conocida está agregada.

Abstract.

This catalogue presents the 40 species of Culicidae (Diptera : Nematocera) reported from Nicaragua. The geographical distribution, synonyms, hosts, diseases transmitted and natural enemies are given for each species. A bibliography of the Nicaraguan species is included.

* Museo Entomológico, A.P. 527, León, Nicaragua.

** Director del Departamento de Entomología Médica del Centro Nacional de Higiene y Epidemiología, Villa Ruben Darío M-254, Managua - 14, Nicaragua.

Introducción.

Los Culicidae forman una familia numerosa de Diptera Nematocera. Las larvas son acuáticas, los adultos pueden ser identificados por la venación alar presentando escamas y la proboscis larga. Son importantes a nivel médico por ser vectores de muchas enfermedades tropicales.


Las larvas de zancudos se encuentran en muchos tipos de aguas, por ejemplo en charcos, huecos o recipientes artificiales, cada especie tiene un tipo de agua característico donde se reproduce. Los huevos son dejados en paquetes sobre la superficie del agua. Las larvas comen algas y materia vegetal en decomposición. Las larvas respiran principalmente a la superficie, ayudándose muchas veces de un sifón. Las pupas son acuáticas y al contrario de los otros insectos, son bastante activas. Respiran en la superficie del agua con dos órganos parecidos a unas trompetitas que poseen sobre el torax. Los adultos se mantienen cerca de su lugar de crianza. Son principalmente nocturnos y descansan durante el día. Las hembras son hematófagas y los machos comen néctar de flores. Se pueden reconocer los machos por las antenas plumosas. Los Culicidae son vectores de enfermedades tales como malaria, dengue, encefalitis, fiebre amarilla, etc. Muchas técnicas de lucha existen ya contra los zancudos, química, ambientales (secado de pantanos) y biológica (bacillus, otros culicidos, peces,...). También la gente puede protegerse con repelentes que se aplican sobre la piel o las ropas.

Especies de Nicaragua.

Aedeomyia squamipennis (LYNCH-ARRIBALZAGA) 1878.

= *Aedes squamipennis* LYNCH-ARRIBALZAGA 1878.

Distribución : Cuba, México, Nicaragua (Chinandega, Rivas), Suramérica.



Aedes aegypti (LINNAEUS), hembra.

Aedes (Stegomyia) aegypti (LINNAEUS) 1762.

yellow fever mosquito, tiger mosquito.

= *Culex aegypti* LINNAEUS 1762.

= *Stegomyia fasciata* THEOBALD.

= *Aedes calopus* MEIGEN.

= *Aedes fasciata* TAYLOR.

= *Aedes mosquito* ROBINEAU-DESVOIDY.

Distribución : Asia, Africa, USA, Cuba, Puerto Rico, Is. Vírgenes, Haití, Trinidad, México,

Guatemala, Honduras, El Salvador, Nicaragua (Chinandega, León, Chontales, Granada, Carazo,

Madriz, Managua, Jinotega, Estelí, Boaco, Nueva Segovia, Matagalpa, Masaya, Rivas,

Zelaya), Panamá.

Hematófago : Mammalia : Hominidae : *Homo*.

Cebidae.

Vector de (fiebre amarilla, yellow fever).

(dengue, break bone fever, quebradora).

(Filariosis humana).

Enemigos naturales :

DIP. Culicidae : *Toxorhynchites brevivalpis*.

Toxorhynchites rutilus.

Bacteria : *Bacillus sphaericus* (Ser.H-5a,5b) (Strain 1593).

Bacillus thuringiensis thuringiensis (Ser.H-1) (Beta-
exotoxina).

Bacillus thuringiensis israelensis (Ser.H-14) (Strain ONR-
60A).

Aedes angustivittatus DYAR & KNAB.

Distribución : Nicaragua (Zelaya).

Aedes atropalpus.

Distribución : Nicaragua (Estelí, León, Managua, Granada, Chontales,

Matagalpa, Zelaya).

Hematófago : Mammifera : Hominidae : *Homo*.

Aedes epaticus DYAR & KNAB.

Distribución : El Salvador, Nicaragua (Matagalpa, Rivas).

Hematófago : Mammifera : Hominidae : *Homo*.

Aedes euplocamus DYAR & KNAB.

Distribución : Nicaragua (Carazo).

Aedes mediovittatus.

Distribución : Nicaragua (Estelí, León, Chinandega, Managua, Masaya,

Granada, Juigalpa,

Matagalpa).

Hematofago : Mammifera : Hominidae : *Homo*.

Aves.

Amphibia : rana, sapo.

Aedes salinarius.

Distribución : Nicaragua (León, Granada).

Hematófago : Mammifera : Hominidae : *Homo*.

Aedes serratus THEOBALD.

Distribución : El Salvador, Nicaragua (Matagalpa).

Hematófago : Mammifera : Hominidae : *Homo*.

Aves.

Reptilia.

Amphibia.

Vector de fiebre de Ilheus.

Aedes sollicitans WALKER.

Distribución : Nicaragua (León, Granada, Matagalpa).

Hematófago : Mammalia : Hominidae : *Homo*.

Vector de encefalitis del este.

dengue.

Aedes (Ochlerotatus) taeniorhynchus (WIEDEMANN) 1821.

black saltmarsh mosquito.

= *Culex taeniorhynchus* WIEDEMANN 1821.

= *Aedes niger* GILES.

= *Aedes portoricensis* LUDL.

= *Aedes damnosus* DYAR & KNAB.

= *Aedes pix* MARTINI 1935.

Distribución : USA, Cuba, Puerto Rico, México, Belize, Nicaragua

(Granada).

Hematófago : Mammalia : Hominidae : *Homo*.

Vector de dengue.

Enemigos naturales :

Fungi : *Culicinomyces clavisporus*.

Bacteria : *Bacillus thuringiensis israelensis* (Ser. H-14).

Aedes terreus WALKER.

Distribución : Nicaragua (Rivas).

Hematófago : Mammalia : Hominidae : *Homo*.

Aedes triseriatus SAY.

Distribución : Nicaragua (Estelí, León, Managua, Granada, Matagalpa).

Hematófago : Mammalia : Hominidae : *Homo*.

Vector de encefalitis californiana.

Anopheles acuasalis CURREY 1942.

= *Anopheles tarsimaculatus* GOELDI 1905.

= *Anopheles emilianus* KOMP 1941.

= *Anopheles delta* ANDUZE 1948.

= *Anopheles guarauno* ANDOZE 1948.

= *Anopheles guaruajensis* RAMOS 1942.

Distribución : Nicaragua (Zelaya, León, Chinandega, Rivas), Ecuador, Brasil,

Uruguay, Argentina.

Hematófago : Mammalia : Hominidae : *Homo*.

Vector de malaria.

filariasis de Bancroft.

Anopheles (Nyssorhynchus) albimanus WIEDEMANN 1821.

= *Anopheles argyrotarsis albipes* THEOBALD 1901.

= *Anopheles cubensis* AGRAMONTE 1900.

= *Nyssorhynchus cubensis*.

= *Anopheles dubius* THEOBALD; BLANCHARD 1905.

= *Anopheles gorgasi* DYAR & KNAB 1907.

= *Anopheles tarsimaculata* GOELDI 1905.

= *Anopheles trisignatus* HOFFMAN 1938.

Distribución : USA, Cuba, Dominica, México, Guatemala, El Salvador,

Honduras,

Nicaragua (Chinandega, León, Rivas, Zelaya), Costa

Rica, Panamá, Ecuador, Brasil.

Hematófago : Mammalia : Hominidae : *Homo*.

Vector de malaria.

Enemigos naturales :

Protozoario : Microsporidia : *Nosema stegomyia* (= *algerae*).

Bacteria : *Bacillus sphaericus* (Ser. H-5a,5b) (Strain 1881-48).

Bacillus thuringiensis israelensis.

Nematoda : Mermithidae : *Reesimermis* (= *Romanomermis*)

culicivorax.

Peces : *Gambusia affinis.*

Poecilia sphenox.

Thilapia sp.

Anopheles apicimacula DYAR & KNAB 1906.

Distribución : Nicaragua (Zelaya), Ecuador, Perú.

Hematófago : Mammalia : Hominidae : *Homo.*

Vector de malaria.

Anopheles argyritarsis ROBINEAU-DESVOIDY 1927.

= *Anopheles rooti* BRETHERS 1926.

Distribución : Nicaragua (Zelaya, Chinandega, Chontales, Managua,

Matagalpa,

Granada), Perú, Bolivia, Brasil, Paraguay, Uruguay,

Argentina.

Hematófago : Mammifera : Hominidae : *Homo.*

Vector de malaria.

Anopheles crucians WIEDEMANN 1828.

Distribución : USA, Caribe, El Salvador, Nicaragua (Zelaya).

Hematófago : Mammalia : Hominidae : *Homo.*

Enemigos naturales :

Nematoda : Mermithidae : *Reesimermis culicivorax*

(=*Romanomermis nielsenii*).

Anopheles darlingi ROOT 1926.

= *Anopheles paulistensis* GALVAO, LANE & CORREA 1937.

Distribución : México, Guatemala, Honduras, Nicaragua (Zelaya), Guyana,

Ecuador,

Perú, Bolivia, Brasil, Paraguay, Argentina, Chile.

Hematófago : Mammalia : Hominidae : *Homo.*

Vector de malaria.

filarias is de Bancroft.

Anopheles eiseni COQUILLET 1902.

= *Anopheles niveopalpis* LUDLOW 1919.

Distribución : México, Nicaragua (Zelaya).

Anopheles hectoris MIRA 1931.

Distribución : El Salvador, Nicaragua (Zelaya).

Hematófago : Mammalia : Hominidae : *Homo*.

Vector de malaria.

Anopheles neivai HOWARD, DYAR & KNAB 1917.

= *Anopheles hylephilus* DYAR & KNAB 1917.

Distribución : Nicaragua (Zelaya).

Anopheles neomaculipalpus CURRY 1931.

Distribución : El Salvador, Nicaragua (Zelaya), Bolivia, Brasil, Paraguay,
Argentina.

Uruguay,

Hematófago : Mammalia : Hominidae : *Homo*.

Anopheles (Anopheles) pseudopunctipennis pseudopunctipennis THEOBALD.

= *Anopheles argentinus* BRETHERS 1912.

= *Anopheles bisignatus* HOFFMANN 1938.

= *Anopheles Perúvianus* TAMAYO 1907.

= *Anopheles tucumanus* LAHILLE 1912.

Distribución : México, Nicaragua (Rivas), Ecuador, Perú, Argentina.

Hematófago : Mammalia : Hominidae : *Homo*.

Vector de Malaria.

Anopheles punctimacula DYAR & KNAB 1906.

= *Anopheles male factor* DYAR & KNAB 1907.

= *Anopheles strigimacula* DYAR & KNAB 1907.

= *Anopheles venezuelae* EVANS 1922.

Distribución : Nicaragua (Zelaya), Costa Rica, Panamá, Colombia, Perú, Bolivia, Brasil,
Uruguay.

Hematófago : Mammalia : Hominidae : *Homo*.

Vector de malaria.

Anopheles strodei ROOT 1926.

= *Anopheles evansae* UNTI 1945.

Distribución : Nicaragua (Zelaya), Bolivia, Brasil, Paraguay, Uruguay, Argentina.

Hematófago : Mammalia : Hominidae : *Homo*.

Vector de malaria.

Anopheles vestitipennis.

Distribución : El Salvador, Nicaragua (Zelaya).

Hematófago : Mammalia : Hominidae : *Homo*.

Culex corniger THEOBALD 1903.

= *Culex lactator* DYAR & KNAB.

= *Culex hassardi* GRABHAM.

= *Culex leucotelus* MAC CORMICK.

Distribución : Cuba, México, Nicaragua (Chinandega, Chontales, Estelí, Matagalpa, Zelaya),
Uruguay.

Culex (Culex) coronator DYAR & KNAB.

Distribución : México, Nicaragua (Chinandega, León, Estelí, Granada, Chontales,
Matagalpa, Carazo, Rivas, Zelaya).

Hematófago : Mammifero : Hominidae : *Homo*.

Culex daumastocampa DYAR & KNAB.

Distribución : Nicaragua (Zelaya).

Culex declarator DYAR & KNAB.

Distribución : Nicaragua (Rivas).

Culex (Melanconion) erraticus DYAR & KNAB.

Distribución : USA, México, Nicaragua (Chinandega, Rivas, Zelaya).

Hematófago : Mamífero : Hominidae : *Homo*.

Culex imitator THEOBALD.

Distribución : Nicaragua (Zelaya).

Culex (Culex) interrogator DYAR & KNAB.

Distribución : México, Nicaragua (Chinandega).

Culex mollis DYAR & KNAB.

Distribución : Nicaragua (Chinandega).

Culex (Culex) nigripalpus THEOBALD 1901.

= *Culex similis* THEOBALD.

= *Culex palus* PALOS.

Distribución : USA, Cuba, México, Nicaragua (Rivas, Estelí, León,

Managua,

Chontales, Matagalpa, Zelaya).

Hematófago : Mammalia : Hominidae : *Homo*.

Enemigos naturales :

Bacteria : *Bacillus sphaericus* (Ser. H-5a-5b) (strain 1593).

Culex pilosus (DYAR & KNAB) 1906.

= *Mochlostyrax pilosus* DYAR & KNAB 1906.

= *Culex ignobilis* DYAR & KNAB.

= *Culex cubensis* DYAR & KNAB.

= *Culex jamaicensis* GRABHAM.

= *Culex floridanus* DYAR & KNAB.

= *Culex agitator* DYAR & KNAB.

Distribución : USA, Cuba, Jamaica, Nicaragua (Zelaya).

Culex (Culex) pipiens quinquefasciatus (SAY) 1828.

southern house mosquito.

= *Culex pipiens fatigans* WIEDEMAN 1828.

= *Culex quinquefasciatus*.

= *Culex fatigans*.

= *Culex cubensis* BIGOT.

Distribución : Asia, Hawaii, USA, Cuba, Puerto Rico, México, Guatemala, El Salvador,
Honduras, Nicaragua (Chinandega, Rivas, Carazo, Zelaya), Costa Rica, Panamá, Colombia,
Venezuela, Guiana francesa, Perú, Brasil.

Hematófago : Mammalia : Hominidae : *Homo*.

Equidae : *Equus*.

Aves (gallinas), *Cyanocitta*, *Mimus*, *Columba*, *Anser*, *Passer*.

Vector de *Brugia sp.*

Dinofilaria immitis (Filariosis canina).

Wuchereria bancrofti (Filariosis de Bancroft).

(Dengue).

(St.Luis encefalitis virus, encefalitis de San Luis).

(Venezuelan equine encephalomyelitis virus).

(Venezuelan equine encephalitis).

Enemigos naturales :

Peces : *Gambusia affinis*.

Nematoda : Mermithidae : *Reesimermis culicivorax*.

Reesimermis nielseni.

Bacteria : *Bacillus thuringiensis israelensis* (Ser.H-14) (Strain ONR-

60A).

Bacillus sphaericus (Ser.H-1a) (Strain SS-II-1 = WHO

1321).

(Ser.H-2) (Strain 1404) (Strain 1894).

(Ser.H-5a,5b) (Strain 1593) (Strain 1881)

(Strain 1691) (Strain 2013-6)

(Strain 2117-2).

(Ser.H-25) (Strain MR-4 = WHO 2297).

Culex restuans THEOBALD.

Distribución : USA, Nicaragua (Estelí, Jinotega, Matagalpa, Chontales,

Zelaya).

Culex salinarius COG.

Distribución : USA, Nicaragua (Granada, Rivas, Zelaya).

Culex tarsalis COG.

Distribución : USA, Nicaragua (Granada, Chontales, Managua, Matagalpa).

Culex (Microculex) sp.

Distribución : Nicaragua (Zelaya).

Culex sp.

Distribución : Nicaragua.

Culiseta sp.

Distribución : Nicaragua (Managua).

Deinocerites epitedeus KNAB.

Distribución : Nicaragua (Chinandega).

Deinocerites pseudus DYAR & KNAB.

Distribución : Nicaragua (Chinandega).

Haemagogus anastasionis DYAR.

= *Haemagogus dominguezi* DURET 1971.

Distribución : Curacao, Nicaragua.

Haemagogus chalcospilans DYAR.

Distribución : Nicaragua (Rivas).

Haemagogus equinus (THEOBALDT).

Distribución : Nicaragua (Rivas).

Limatus durhami THEOBALD 1901.

Distribución : Cuba, Trinidad, El Salvador, Nicaragua (Zelaya), Costa Rica, Panamá,
Colombia, Venezuela, Guyana, Ecuador, Perú, Bolivia, Brasil.

Megarhinus moctezuma DYAR & KNAB.

Distribución : Nicaragua (Chinandega).

Orthopodomyia kummi EDWARDS.

Distribución : México, Nicaragua (Chinandega).

Phoniomyia lassalli (BONNE-WEPSTER & BONNE).

Distribución : Nicaragua (Zelaya).

Psorophora (Grabhamia) confinnis (LYNCH ARRIBALZAGA) 1891.

florida glade mosquito.

= *Taeniorhynchus confinnis* LYNCH ARRIBALZAGA 1891.

= *Psorophora jamaicensis* THEOBALD.

Distribución : USA, Cuba, Jamaica, México, El Salvador, Nicaragua (Granada,

Estelí, Managua, Zelaya, Nueva Segovia, Chinandega, León, Granada, Masaya, Carazo, Chontales,

Boaco, Matagalpa, Rivas).

Hematófago : Mammalia : Hominidae : *Homo*.

Psorophora ferox (HUMBOLDT) 1820.

= *Culex ferox* HUMBOLDT 1820.

= *Aedes sayi* DYAR & KNAB.

= *Aedes pazozi* DYAR & KNAB.

Distribución : USA, Cuba, Nicaragua (Chinandega).

Hematófago : Mammalia : Hominidae : *Homo*.

Toxorhynchites (Lynchiella) rutilus rutilus (COQUILLET).

Distribución : El Salvador, Nicaragua (Estelí, León, Managua, Granada).

Depredador de DIP. Culicidae : *Aedes aegypti*.

Uranotaenia geometrica THEOBALD.

Distribución : Nicaragua (Zelaya).

Uranotaenia lowii THEOBALD 1901.

= *Uranotaenia continentalensis* DYAR & KNAB.

= *Uranotaenia minuta* THEOBALD.

= *Uranotaenia monilis* SH. & P.

Distribución : Cuba, Nicaragua (Rivas).

Wyeomyia celaenocephala (DYAR & KNAB).

Distribución : Nicaragua (Zelaya).

Wyeomyia circumcincta (DYAR & KNAB).

Distribución : Nicaragua (Zelaya).

Bibliografía.

ACREE F., TURNER R.B., GOUCK H.K., BEROZA M. & SMITH N. (1968) L-lactic acid: a mosquito attractant isolated from humans. *Science*, 161:1346-1347.

AITKEN T.H.G. (1942) Contributions toward a knowledge of the insect fauna of Lower California. No.6: Diptera: Culicidae. *Proc. Calif. Acad. Sci. Fourth Series*, 24:161-170.

ALAYO-DALMAU P. & GARCIA-AVILA I. (1983) Lista anotada de los dípteros de Cuba. Ed. Científico-Técnica, Habana, Cuba, 201pp.

ALTMAN R.M. (1969) Repellent tests against *Anopheles albimanus* Wiedemann in the Panamá Canal Zone. *Mosq. News*, 29:110-112.

Anónimo (1970) Reinfestation by *Aedes aegypti* in México. *WHO Wkly. Epidemiol. Rep.*, 42:223.

Anónimo (1971) Reinfestation by *Aedes aegypti* in Costa Rica. *WHO Wkly. Epidemiol. Rep.*, 43:95.

Anónimo (1974) Vigilancia de la encefalitis en las Américas. Noviembre. *Vigilancia Epidemiológica, Of. Sanit. Panam.*, 5pp.

Anónimo (1983) Standard method of field testing topical application of compounds as repellents for medically important and pest arthropods (including insects, ticks, and mites): I. Mosquitoes. *Am.Soc. Testing Materials*, E939-83. In *Annual book of ASTM standards*, Vol.11(04):5pp.

ARNELL J.H. (1973) Mosquito studies (Diptera: Culicidae). XXXII. A revision of the genus *Haemagogus*. *Contrib. Am. Ent. Inst. (Ann Arbor)*, 10(2):1-174.

BACOT A. & TALBOT G. (1919) The comparative effectiveness of certain culicifuges under laboratory conditions. *Parasitology*, 11:221-236.

BARJAC H.De (1978) Toxicite de *Bacillus thuringiensis* var. *israelensis* pour les larves d'*Aedes aegypti* et d'*Anopheles*

stephensi. C. R. Acad. Sci. (Paris), 286D:1175-1178.

BARJAC H.De (1978) Un nouveau candidat a la lutte biologique contre les moustiques: *Bacillus thuringiensis* var. *israelensis*. Entomophaga, 23(4):309-319.

BARJAC H.De & LARGET-THIERY I. (1984) Characteristics of IPS-82 as standard for biological assay of *Bacillus thuringiensis* H-14 preparations. WHO/VBC/84.892:10pp.

BARJAC H.De, LARGET-THIERY I., COSMAO-DUMANOIR V. & RIPOUTEAU H. (1985) Serological classification of *Bacillus sphaericus* strains in relation with toxicity to mosquito larvae. Appl. Microbiol. Biotechnol., 21:85-90.

BARR A.R. (1957) The distribution of *Culex p.pipiens* and *C.quinquefasciatus* in North America. Am. J. Trop. Med., 6:153.

BARR A.R., KNIGHT K.L., PILLAI J.S., RAI K.S., ROBBINS W.E. & WARD R.A. (1973) Mosquitoes control. Some perspectives for developing countries. Nat. Acad. Sciences, 63pp.

BAR-ZEEV M. & BEN-TAMAR D. (1971) Evaluation of mosquito repellents. Mosq. News, 31(1):56-61.

BAR-ZEEV M., MAIBACH H.I. & LHAN A.A. (1977) Studies on the attraction of *Aedes aegypti* (Diptera: Culicidae) to man. J. Med. Ent., 14:113-120.

BAUER P., RUDIN W. & HECKER H. (1977) Ultrastructure changes in midgut cells of female *Aedes aegypti* L. (Insecta, Diptera) after starvation or sugar diet. Cell Tissue Research, 177:215-219.

BELKIN J.N. (1977) *Aedes (Ochlerotatus) pix* Martini 1935 a synonym of *Aedes (O.) taeniorhynchus* (Wiedeman 1821). Mosq. Syst., 9:525.

BERNER L. (1947) Notes on the breeding habitats of *Aedes (Stegomyia) aegypti* (Linnaeus). Ann. Ent. Soc. Am., XL(3):528-529.

BLANCHARD (1905) Les moustiques, Histoire naturelle et medicale. 673pp.

BORROR D.J., DE LONG D.M. & TRIPLEHORN C.A. (1981) An introduction to the study of insects. Ed.V. Saunders, USA, 827pp.

BOURGOUIN C., TINELLI R., BOUVET J.P. & PIRES R. (1984) *Bacillus sphaericus*: 1593-4 purification of fraction toxic for mosquito larvae. In Bacterial Protein Toxins, pp.337-338.

BREELAND S.G. (1980) A bibliography to the litterature of *Anopheles albimanus* (Diptera: Culicidae). Mosq. Syst., 12:50-150.

BROOKMAN B. & REEVES W.C. (1953) New records of mosquitoes from Lower California, México, with notes and descriptions. (Diptera: Culicidae). Ann. Ent. Soc. Amer., 46:225-236.

BUESCHER M.D., RUTLEDGE L.C. & WIRTZ R.A. (1982) Tests of commercial repellents on human skin against *Aedes aegypti*. Mosq. News, 42:428-433.

CHARLES J.F. & BARJAC H.De (1981) Histopathologie de l'action de la delta-endotoxine de *Bacillus thuringiensis* var.*israelensis* sur les larves d'*Aedes aegypti* (Dipt.Culicidae). Entomophaga, 26:203-212.

CHARLES J.F. & BARJAC H.De (1983) Action des cristaux de *Bacilius thuringiensis* var. *israelensis* sur l'intestin

moyen de larves de *Aedes aegypti* L., en microscopie electronique. Ann. Microbiol., 134A:197-218.

CHARLES J.F. (1983) Action de la delta-endotoxine de *Bacillus thuringiensis* var. *israelensis* sur cultures de cellules de *Aedes aegypti* L. Ann. Microbiol., 134A:365-381.

CHARLES J.F. (1984) *Bacillus thuringiensis*: crystal biogenesis and cytopathology on mosquito larvae. In Bacterial protein toxins, pp.383-384.

CHARLES L.J. & SENEVET G. (1953) The distribution of *Anopheles albimanus* in the caribbean islands. Am. J. Trop. Med. Hyg., 2:1109-1115.

CHRISTOPHERS S.R. (1960) *Aedes aegypti* (L.) the yellow fever mosquito, its life history, bionomics and structure. Cambridge Univ. Press., London.

COLLINS W.E. (1963) Transmission of St.Louis encephalitis virus by larval-infected *Culex quinquefasciatus* mosquitoes. Ann. Ent. Soc. Am., 56:237-239.

CORDELLIER R., BOUCHITE B., ROCHE J.C. & AKOLIBA P. (1982) Enquete entomologique et epidemiologique sur un cas mortel de fièvre jaune survenu dans la sous prefecture d'Anayama en republique de Cote d'Ivoire. Med. Trop., 42(3):269-273.

CROVELLO T.J. & HACKER C.S. (1972) Evolutionary strategies in life table characteristics among feral and urban strains of *Aedes aegypti* (L.). Evolution, 26:185-196.

DAVIDSON E.W., SINGER S.S. & BRIGGS J.D. (1975) Pathogenesis of *Bacillus sphaericus* strain SS-II-1. Infections of *Culex pipiens quinquefasciatus* (= *C.pipiens fatigans*) larvae. J. Invertebr. Pathol., 25:179-184.

DE MEILLON B., SEBASTIAN A. & KHAN Z. (1967) The duration of egg, larval and pupal stages of *Culex pipiens fatigans* in Rangoon, Burma. Bull. WHO, 36:7-14.

DE MEILLON B., GRAB B. & SEBASTIAN A. (1967) Evaluation of *Wucheria bancrofti* infection in *Culex pipiens fatigans* in Rangoon, Burma. Bull. WHO, 36:91-100.

DESOWITZ R.S. & CHELLAPAH W.T. (1962) The transmission of *Brugia sp.* through *Culex pipiens fatigans*: the effect of age and prior non-infective blood on the infection rate. Trans. Roy. Soc. Trop. Med. Hyg., 56:121.

DES ROCHERS B. & GARCIA R. (1983) The efficacy of *Bacillus sphaericus* in controlling mosquitoes breeding in sewer effluent. Proc. Pap. 51th Ann. Conf. Calif. Mosq. Vect. Control Ass., pp.35-37.

DES ROCHERS B. & GARCIA R. (1984) Evidence for persistance and recycling of *Bacillus sphaericus*. Mosq. News, 44(2):160-165.

DIAZ NAJERA A. (1967) Datos entomológicos obtenidos durante el desarrollo del programa para erradicar *Aedes aegypti* de la República Mexicana. Rev. Invest. Salud Pública, 27:225-232.

DIAZ-NAJERA A. & VARGAS L. (1973) Mosquitoes mexicanos. Distribución geografica actualizada. Rev. Inv. Salud Pública (México), 33:111-125.

- DOWNS W.G. & BORDAS E.** (1949) A malaria survey of the Southern Territory of Lower California. *Amer. J. Trop. Med.*, 29:695-699.
- DURET J.P.** (1971) Una nueva especie de *Haemagogus* de Nicaragua (Diptera: Culicidae). *Neotropica*, 17:83-86.
- DYAR & KNAB** (1917) The mosquitoes of North and Central America and the West Indies. Systematic description part II. Carnegie Inst. Wash., Publ. 159. 4:525-1064.
- FARAN M.E.** (1980) Mosquito studies (Diptera, Culicidae) XXXIV. A revision of the *albimanus* section of the subgenus *Nyssorhynchus* of *Anopheles*. *Contrib. Am. Ent. Inst. (Ann Arbor)*, 15(7):1-215.
- FOX D.A., SEAWRIGHT J.A. & HALL D.W.** (1979) Field survival, migration and ovipositional characteristics of laboratory-reared *Toxorhynchites rutilus rutilus* (Diptera: Culicidae). *J. Med. Ent.*, 16(2):121-127.
- FOX D.A. & BOSTON M.D.** (1979) A quantified mass-rearing technique for *Toxorhynchites rutilus rutilus* Coquillet. *Mosq. News*, 39(3):616-619.
- FOX I. & GARCIA-MOLL I.** (1961) Multi-resistant *Aedes aegypti* in Puerto Rico and Virgin Islands. *Science*, 133:646-647.
- FRAZIER C.A.** (1969) Insect allergy. Warren H.Green, Inc., USA, 493pp.
- GALINDO P. & TRAPIDO H.** (1957) Forest mosquitoes associated with sylvan yellow fever in Nicaragua. *Am. J. Trop. Med. Hyg.*, 6:145-152.
- GARCIA R.** (1983) Mosquito management: ecological approaches. *Env. Manag.*, 7(1):73-78.
- GARCIA R. & SWEENEY A.W.** (1986) The use of microbial pathogens for the control of mosquitoes. *Agric. Ecosyst. and Envir.*, 15:201-208.
- GEERING K. & FREYVOGEL T.A.** (1974) The distribution of acetylcholine and unspecific esterases in the midgut of female *Aedes aegypti* L. *Comp. Biochem. Physiol.*, 49B:775-784.
- GOELDI** (1905) Os mosquitos no Para. Memorias do Museu Goeldi (Museu Paranense), 4:154pp.
- GILBERT I.H., GOUCK H.K. & SMITH C.N.** (1966) Attractiveness of men and women to *Aedes aegypti* and relative protection time obtained with deet. *Fla. Entomol.*, 49(1):53-66.
- GOLDBERG L. & MARGALIT J.** (1977) Bacterial spore demonstrating rapid larvicidal activity against *Anopheles sergentii*, *Uranotaenia unguiculata*, *Culex univittatus*, *Aedes aegypti* and *Culex pipiens*. *Mosquito News*, 37:355-358.
- GOLDBERG L.J., GOLDBERG E.M. & MARGALIT J.** (1977) Potential application of a bacterial spore, ONR-60A to mosquito larval control: demonstrated rapid larvicidal activity against *Anopheles sergentii*, *Uranotaenia unguiculata*, *Culex univittatus*, *Aedes aegypti* and *Culex pipiens* (complex). WHO/VBC/77.662.
- GOMEZ C., RABINOVICH J.E. & MACHADO-ALLISON C.E.** (1977) Population analysis of *Culex pipiens fatigans* Wied. (Diptera: Culicidae) under laboratory conditions. *J. Med. Ent.*, 13(4):453-463.
- GONZALEZ-BROCHE R.** (1985) Nuevos reportes sobre la tribu Sabethini (Diptera: Culicidae) para Cuba. *Poeyana*, 298:11pp.
- GOUCK H.K., GODWIN D.R., SCHRECK C.E. & SMITH N.** (1967) Field tests with repellent-treated netting against black salt-marsh mosquitoes. *J. Econ. Ent.*, 60(5):1451-1452.

- GOUCK H.K., MAC GOVERN T.P. & BEROZA M.** (1967) Chemical tested as space repellent against yellow-fever mosquitoes. *J. Econ. Ent.*, 60(6):1587-1590.
- HAMMON W. & REEVES W.C.** (1943) Laboratory transmission of St.Louis encephalitis virus by three genera of mosquitoes. *J. Exp. Med.*, 78:241.
- HARBACH R.E., DAHL C. & WHITE G.B.** (1985) *Culex (Culex) pipiens* LINNAEUS (Diptera: Culicidae): concepts, type designations, and description. *Proc. Ent. Soc. Wash.*, 87(1):1-24.
- HEINEMANN S.J. & BELKIN J.N.** (1977) Collection records of the project "Mosquitoes of Middle America" 8. Central America: Belize(BH); Guatemala(GUA), El Salvador(SAL), Honduras(HON), Nicaragua(NI,NIC). *Mosq. Syst.*, 9:403-454.
- HERTLEIN B.C., LEVY R. & MILLER T.W.** (1979) Recycling potential and selective retrieval of *Bacillus sphaericus* from soil in a mosquito habitat. *J. Invert. Pathol.*, 33:217-221.
- HORNBY J.A., HERTLEIN B.C., LEVY R. & MILLER T.W.** (1981) Persistent activity of mosquito larvicidal *Bacillus sphaericus* 1593 in fresh water and sewage. WHO/VBL/81.830.
- IGNOFFO C.M., COUCH T.L., GARCIA C. & KROHA M.J.** (1980) Relative activity of *Bacillus thuringiensis* var. *kurstaki* and var. *israelensis* against larvae of *Aedes aegypti* (L.), *Culex pipiens quinquefasciatus* Say, *Trichoplusia ni* (Hubner), *Heliothis zea* (Boddie) and *Heliothis virescens* (F.). *J. Econ. Ent.*
- JARRETT P.** (1985) Potency factors in the delta-endotoxin of *Bacillus thuringiensis* var. *aizawi* and the significance of plasmids in their control. *Jl. Appl. Bacteriol.*, 58:437-448.
- JORDAN P.** (1962) Transmission of *Brugia* through *Culex pipiens fatigans*. *Trans. Roy. Soc. Trop. Med. Hyg.*, 56:436.
- KELLOG F.E., BURTON D.J. & WRIGHT R.H.** (1968) Measuring mosquito repellency. *Can. Ent.*, 100:763-768.
- KHAN A.A., MAIBACH H.I., STRAUSS W.G. & FENLEY W.R.** (1966) Quantation of effects of several stimuli on the approach of *Aedes aegypti*. *J. Econ. Ent.*, 59(3):690-694.
- KHAN A.A. & MAIBACH H.I.** (1966) Quantation of effect of several stimuli on landing and probing by *Aedes aegypti*. *J. Econ. Ent.*, 59(4):902-905.
- KHAN A.A. & MAIBACH H.I.** (1972) Insect repellents. 1. Effect on the flight and approach by *Aedes aegypti*. *J. Econ. Ent.*, 65(5):1318-1321.
- KITZMILLER J.B.** (1982) Anopheline names: their derivations and histories. *Ent. Soc. Amer.*, Thomas Say Foundation, VIII:639pp.
- KNIGHT K.L. & STONE A.** (1977) A catalog of the mosquitoes of the world (Diptera: Culicidae). *Amer. Ent. Soc.*, Thomas Say Foundation, VI:611pp.
- KNIGHT K.L.** (1978) Supplement to a catalog of the mosquitoes of the world (Diptera: Culicidae). *Amer. Ent. Soc.*, Thomas Say Foundation, VI(Suppl.):107pp.
- KOMP W.H.W.** (1941) The classification and identification of the *Anopheles* mosquitoes of México, Central America and the West Indies. *Amer. Assoc. Adv. Sci. Publ.*, 15:88-97.

- KOMP W.H.W.** (1942) The anopheline mosquitoes of the Caribbean region. U.S. Publ. Health Serv., Natl. Inst. Health Bull., 179:195pp.
- KOST A.N., TEREN'TEV P.B., ELIZAROV, YU A. & TSYBA I.F.** (1971) Perception of organic compounds by *Aedes aegypti* mosquitoes. Khernmoresteptsiya Nasekomykh. Mater. Vses. Simp. 1st, pp.89-94.
- KRISHNAMURTI B.S. & PAL R.** (1958) A note on the comparative effect of different blood feed on production of *Culex pipiens fatigans*. Bull. Nation. Soc. Ind. Malar. Mosq. Dis., 17:27.
- KUMM H.W.** (1942) *Anopheles crucians* found in Northern Nicaragua. Amer. Jour. Trop. Med., 22(5):511-512.
- LAHKIM-TSOR L., PASCAR-GLUZMAN C., MARGALIT J. & BARAK Z.** (1983) Larvicidal activity of *Bacillus thuringiensis* subsp. *israelensis*, serovar H 14 in *Aedes aegypti*: histopathological studies. J. Invert. Path., 41:104-116.
- LANE J.** (1953) Neotropical Culicidae. Univ. Press, Sao Paulo, 1112pp.
- LARGET-THIERY I. & BARJAC H.De** (1981) Specificite et principe actif de *Bacillus thuringiensis* var. *israelensis*. Bull. Soc. Pathol. Exot., 74(2):216-227.
- LARGET-THIERY I., HAMON S. & BARJAC H.De** (1984) Sensibilite des Culicidae a la beta-exotoxine de *Bacillus thuringiensis*. Entomophaga, 29(1):95-108.
- LAURENT P. & CHARLES J.F.** (1984) Action comparee des cristaux solubilises des serotypes H-14 et H-1 de *Bacillus thuringiensis* sur des cultures de cellules de *Aedes aegypti*. Ann. Microbiol., 135A:473-484.
- LAVEN H.** (1967) Speciation and evolution in *Culex pipiens*. In WRIGHT J.W. & PAL R. Genetics of insect vectors of disease. Elsevier, Amsterdam, pp.251-275.
- LAVEN H.** (1967) Eradication of *Culex pipiens fatigans* through cytoplasmic incompatibility. Nature, 216:383.
- LEE R.** (1974) Structure and function of the fascicular stylets, and labral and cibarial sense organs of male and female *Aedes aegypti* (L.) (Diptera, Culicidae). Quaest. Ent., 10:187-215.
- LOPEZ C.** (1972) Efectos de la densidad larval sobre componentes de las tabla de vida en *Aedes aegypti* L. (Dipt. Culicidae). Thesis, School Biol., Central Univ. Venezuela.
- MACHADO-ALLISON C.E.** (1971) Genetic differences among subspecies of *Aedes aegypti* (L.). Ph.D. Thesis, Univ. Notre Dame, 158pp.
- MARTINEZ-PALACIOS A.** (1986) Notas sobre los mosquitos de Baja California, México (Diptera: Culicidae). Fol. Ent. Mex., 69:99-116.
- MARTINI E.** (1935) Los mosquitos de México. Dept. Salubr. Publ., Bol. Tecn., Ser.A, Ent. Med. Parasit., 1:66pp.
- MATTINGLY P.E.** (1962) Some considerations relating to the role of *Culex pipiens fatigans* in the transmission of human filariasis. Bull. Wld. Hlth. Organ., 27:569.
- MAYER M.S. & JAMES J.D.** (1969) Attraction of *Aedes aegypti*: responses to human harms, carbon dioxide and air currents in a new type of olfactometer. Bull. Ent. Res., 58(3):629-642.
- MOUNT G.A., HAILE D.C., PIERCE N.W. & BALDWIN K.R.** (1978) Aerosol of permethrin and propuxur for control of adult mosquitoes. Mosq. News, 38(1):39-42.
- MYERS P. & YOUSTEN A.** (1978) Toxic activity of *Bacillus sphaericus*, SSII-1 for mosquito larvae. Infect. Immun.,

19:1047-1053.

NAYAR J.K. (1968) Biology of *Culex nigripalpus* Theobald (Diptera: Culicidae). Part 1: Effects of rearing conditions on growth and the diurnal rhythm of pupation and emergence. J. Med. Ent., 5:39-46.

NAYAR J.K. & SAVERMAN D.M. (1970) A comparative study of growth and development in Florida mosquitoes. Part 1: Effects of environmental factors on ontogenic timings, endogenous diurnal rhythm and synchrony of pupation and emergence. J. Med. Ent., 7:163-174.

NICKLE W.R. (1979) Probable establishment and overwintering of a Mermithid nematode parasite of mosquitoes in Maryland. Proc. Helminthol. Soc. Wash., 46(1):21-27.

NIELSEN E.T. & NIELSEN A.T. (1953) Field observations on habits of *A. taeniorhynchus*. Ecology, 34:141.

PATTERSON R.S., WEIDHAAS D.E., FORD H.R. & LOFGREN C.S. (1970) Suppression and elimination of an island population of *Culex pipiens quinquefasciatus*. Science, 168:1368-1370.

PERRY W.J. (1950) The dengue vector on New Caledonia, the New Hebrides and the Solomon islands. Am. J. Trop. Med., 28(2):253-259.

PETERSON J.J., WILLIS O.R. & CHAPMAN H.C. (1978) Release of *Romanomermis culicivorax* for the control of *Anopheles albimanus* in El Salvador.I. Am. J. Trop. Med. Hyg., 27:1265-1266.

PETERSON J.J., CHAPMAN H.C., WILLIS O.R. & FUKADA T. (1978) Release of *Romanomermis culicivorax* for the control of *Anopheles albimanus* in El Salvador.II. Am. J. Trop. Med. Hyg., 27:1268-1273.

PLETSCH D.J. (1979) Initial records of *Anopheles albimanus* in the peninsula of Baja California, México. Mosq. News, 39:799-800.

PROVOST M.W. (1951) Occurrence of saltmarsh mosquitoes in interior of Florida. Fla. Ent., 34:48.

PURCELL B.H. (1981) Effects of *Bacillus thuringiensis* var. *israelensis* on *Aedes taeniorhynchus* and some non-target organisms in the salt marsh. Mosq. News, 41(3):476-483.

QUTUBUDDIN M. (1952) The emergence and sex ratio of *Culex pipiens fatigans* Wied. (Diptera, Culicidae) in laboratory experiments. Bull. Ent. Res., 43:549.

RAGEAU J. (1955) Insectes et autres arthropodes d'interet medical ou veterinaire en Nouvelle Caledonie et aux iles Loyaute. Etudes Melanesiennes, pp.60-104.

RAI K.S. & MAC DONALD P.T. (1971) Chromosomal translocation and genetic control of *Aedes aegypti*. In The sterility principles for insect control or eradication. Int. Atomic Energy Agency Press, Vienna, Austria, pp.437-452.

RICHARDS O.W. & DAVIES R.G. (1977) Imm's general textbook of entomology. Ed.X. Chapman & Hall, London, Vol.II:421-1354.

ROESSLER P. (1963) The attractiveness of steroids and amino acids to female *Aedes aegypti*. Proc. 50th Ann. Meeting New Jersey Mosq. Extermination Assoc. and 19th Ann. Meeting Amer. Mosq. Control Assoc.

ROSS H.H. & HORSFALL W.R. (1965) A synopsis of the mosquitoes of Illinois (Diptera, Culicidae). Illinois Nat. Hist. Surv., Biol. Notes, 52:50pp.

- ROY D.W. & MAJUMDAR S.P.** (1939) On mating and egg formation in *Culex pipiens fatigans* Wied. J. Malar. Inst. India, 2:245.
- ROZEBOOM L.E.** (1953) A note on the presence of *Anopheles albimanus* in Barbados. Am. J. Trop. Med. Hyg., 2:1116.
- RUBIO-MORAN R., ROWLEY W.A. & COATS J.R.** (1981) Effect of permethrin and malathion on *Aedes trivittatus* and *Culex pipiens pipiens*. Iowa Ste. Jl. Res., 55(3):235-244.
- RUDIN W. & HECKER H.** (1976) Morphometric comparison of the midgut epithelial cells in male and female *Aedes aegypti* L. (Insecta: Diptera). Tissue and Cell, 8:456-470.
- SAN MARTIN-BARBIERI C., GROOT N. & OSORNO-MESA E.** (1953) Human epidemic in Colombia caused by Venezuelan equine encephalomyelitis virus. Am. J. Trop. Med. Hyg., 3:283.
- SCHRECK C.E. & JAMES J.** (1968) Broth cultures of *Bacteria* that attract female mosquitoes. Mosq. News, 28(1):33-38.
- SCHRECK C.E., GOUCK H.K. & POSEY K.H.** (1970) An experimental plexiglas mosquito trap utilizing carbon dioxide. Mosquito News, 30(4):641-645.
- SCHRECK C.E., GILBERT I.H., WEIDHAAS D.E. & POSEY K.H.** (1970) Spatial action of mosquito repellents. J. Econ. Ent., 63(5):1576-1578.
- SCHRECK C.E., GOUCK H.K. & POSEY K.H.** (1972) The range of effectiveness and trapping efficiency of a plexiglass mosquito trap baited with carbon dioxide. Mosquito News, 32(4):496-501.
- SCHRECK C.E., WEIDHAAS D.E., SMITH N. & POSEY K.H.** (1977) Chemical treatment of wide-mesh net clothing for personal protection against blood-feeding arthropods. Mosq. News, 37(3):455-462.
- SCHRECK C.E. & SMITH N.** (1977) Repellency of cyclohexanethylene carbamide and N,N-diethyl-m-toluamide against *Aedes aegypti* and *Ae. taeniorhynchus*. Mosq. News, 37(3):525-527.
- SCHRECK C.E., WEIDHAAS D.E. & SMITH N.** (1977) Evaluation of electronic sound-producing devices against *Aedes taeniorhynchus* and *Ae. sollicitans*. Mosq. News, 37(3):529-531.
- SCHRECK C.E.** (1977) Techniques for the evaluation of insect repellents: a critical review. Ann. Rev. Ent., 22:101-119.
- SCHRECK C.E., POSEY K. & SMITH D.** (1978) Durability of permethrin as a potential clothing treatment to protect against blood-feeding arthropods. J. Econ. Ent., 71(3):397-400.
- SCHRECK C.E., SMITH N., WEIDHAAS D., POSEY K. & SMITH D.** (1978) Repellent vs. toxicants as clothing treatments for protection from mosquitoes and other biting flies. J. Econ. Ent., 71(6):919-922.
- SCHRECK C.E., CARLSON D.A., WEIDHAAS D.E., POSEY K. & SMITH D.** (1980) Wear and aging tests with permethrin-treated cotton-polyester fabric. J. Econ. Ent., 73(3):451-453.
- SCHRECK C.E., SMITH N., CARLSON D.A., PRICE G.D., HAILE D. & GODWIN D.R.** (1981) A material isolated from human hands that attracts female mosquitoes. Jl. Chemical Ecol., 8(2):429-438.
- SCHRECK C.E., KANE F. & CARLSON D.A.** (1982) Permethrin impregnations of military fabrics: an evaluation of application rates and industrial methods by bioassay and gas chromatographic analysis. Soap, Cosmetic, Chemical specialities, pp.36-39.

- SCHRECK C.E., KLINE D.L., CHANIOTIS B.N., WILKINSON N., MAC GOVERN T.P. & WEIDHAAS D.E.** (1982) Evaluation of personal protection methods against phlebotomine sand flies including vectors of leishmaniasis in Panamá. *Am. J. Trop. Med. Hyg.*, 31(5):1046-1053.
- SCHRECK C.E., WEBB J.C. & BURDEN G.S.** (1984) Ultrasonic devices: evaluation of repellency to cockroaches and mosquitoes and measurement of sound output. *J. Environ. Sci. Health*, A19(5):521-531.
- SCHRECK C.E., HAILE D.G. & KLINE D.L.** (1984) The effectiveness of permethrin and deet, alone or in combination, for protection against *Aedes taeniorhynchus*. *Am. J. Trop. Med. Hyg.*, 33(4):725-730.
- SCHRECK C.E.** (1985) The status of deet (N,N-Diethyl-M-Toluamide) as a repellent for *Anopheles albimanus*. *J. Am. Mosq. Control Ass.*, 1(1):98-100.
- SCHRECK C.E. & MAC GOVERN T.P.** (1985) Repellent tests in the field and laboratory against wild populations of *Mansonia titillans* (Diptera: Culicidae). *J. Med. Ent.*, 22(6):658-662.
- SCORZA J.V.** (1972) Observaciones bionómicas sobre *Culex pipiens fatigans* Wied. 1829. Ediciones del Rectorado, Univ. de los Andes, Merida, 198pp.
- SHRIVER D. & BICKLEY W.E.** (1964) The effect of temperature on hatching of eggs of the mosquito, *Culex pipiens quinquefasciatus* Say. *Mosquito News*, 24:137-140.
- SINGH K.R.P.** (1967) Cell cultures derived from larvae of *Aedes albopictus* (Skuse) and *Aedes aegypti* (L.). *Curr. Sci.*, 36:506-508.
- SMITH C.N., SMITH N., GOUCK H.K., WEIDHAAS D.E., GILBERT I.H., MAYER M.S., SMITTLE B.J. & HOFBAUER A.** (1970) L-lactic acid as a factor in the attraction of *Aedes aegypti* (Diptera: Culicidae) to human hosts. *Ann. Ent. Soc. Am.*, 63(3):760-770, 65:607-612.
- SMITH C. N., SCHRECK C.E. & SMITH D.** (1976) Field studies with selected skin repellents against natural infestations of *Aedes taeniorhynchus* (Wiedemann). *Mosq. News*, 36:36-38.
- SOUTHWOOD T.R.E., MURDIE G., YASUNO M., TONN R. & READER P.** (1972) Studies on the life of *Aedes aegypti* in Wat Samphaya, Bangkok, Thailand. *Bull. WHO*, 46:211-226.
- STEFFAN W.A.** (1981) The type-specimens of *Toxorhynchites* (Diptera: Culicidae) at the National Museum of Natural History. *Mosq. News*, 12:379-385.
- SUAREZ O. & BERGOLD G.H.** (1968) Investigations on an outbreak of Venezuelan equine encephalitis in towns of eastern Venezuela. *Amer. J. Trop. Med. Hyg.*, 17:875.
- TAYLOR R.T. & SCHOOF H.F.** (1971) The relative effectiveness of malathion thermal aerosol and ground-applied ULV against three species of mosquitoes. *Mosq. News*, 31(3):346-349.
- TAYLOR W.G. & SCHRECK C.E.** (1985) Chiral-phase capillary gas chromatography and mosquito repellent activity of some oxazolidine derivatives of (+)- and (-)-citronellal. *Jl. Pharm. Sci.*, 74(5):6pp.
- THEOBALD** (1901) A monograph of the Culicidae or mosquitoes. 1:125-128, 2:322-339.
- THEOBALD** (1903) A Monograph of the Culicidae or mosquitoes. 3:173.

- TRPIS M.** (1972) Interaction between the predator *Toxorhynchites brevivalpis* and its prey *Aedes aegypti*. Bull. WHO, 49:359-365.
- TRPIS M.** (1973) Predator-prey oscillations in density of *Toxorhynchites brevivalpis* and *Aedes aegypti* in a suburban habitat in East Africa. Bull. WHO.
- VAN DER KUYP E.** (1949) Notes on *Haematogogus anastasionis* Dyar of Curacao. Doc. Neerl. Indones. Morbios. Trop., 1:142-144.
- VLADIMIROVA V.V.** (1969) Sensitivity of *Aedes aegypti* mosquitoes to repellents depending upon environment. Med. Parazit. Bolezn., 38(2):214-218.
- WARD A.** (1982) Culicidae. In HURLBERT S.H. & VILLALOBOS-FIGUERO A. Aquatic biota of México, Central America and the West Indies. San Diego Ste Univ., California, pp.417-429.
- WARD R.A.** (1984) Second supplement to "A catalog of the mosquitoes of the world" (Diptera: Culicidae). Mosq. Syst., 16(3):227-270.
- WATTAL B.L., KALRA N.L. & BEDI K.M.** (1961) Studies on culicine mosquitoes. 2.Laboratory studies on the longevity of adult *Culex pipiens fatigans* Wied. Indian Jl. Malar., 4:321.
- WATTAL B.L. & KALRA N.L.** (1963) Studies on culicine mosquitoes. 3.Laboratory studies on the effect of the blood of different vertebrates egg production and biology of *Culex fatigans* Wied. Indian Jl. Malar., 17:75.
- WICKREMENSINGHE R.S.B. & MENDIS C.L.** (1980) *Bacillus sphaericus* spore from Sri Lanka demonstrating rapid larvicidal activity on *Culex quinquefasciatus*. Mosq. News, 40:387-389.
- WILLIS O.R., CHAPMAN H.C. & PETERSON J.J.** (1980) Additional field testing of the mermithid parasite *Romanomermis culicivorax* against *Anopheles albimanus* in El Salvador. Mosq. News, 40:71-72.
- WIRTZ R.A., TURRENTINE J.D. & RUTLEDGE L.C.** (1980) Mosquito area repellents: laboratory testing of candidate materials against *Aedes aegypti*. Mosq. News, 40:432-439.
- WOKE P.A.** (1947) Arthropods of sanitary importance in the republic of Nicaragua, Central America. Am. J. Trop. Med., 27:357-375.
- WOOD P.W.** (1968) The effect of ambient humidity on the repellency of ethyl-hexamediol (6'-12) to *Aedes aegypti*. Can. Ent., 100:1331-1335.
- WRIGHT R.H. & BURTON D.J.** (1969) Pyrethrum as a repellent. Pyrethrum Post, 10(2):14-21.
- WRIGHT R.H.** (1975) Why mosquito repellent repel. Sci. Am., 1(4):1-12.
- YOUSTEN A.A., BARJAC H.De, HENDRICK J., COSMAO-DUMANOIR V. & MYERS P.** (1980) Comparison between bacteriophage typing and serotyping for the differentiation of *Bacillus sphaericus* strains. Ann. Microbiol., 131B:297-308.