

chemicalWATCH Factsheet

MALATHION

What is Malathion?

Malathion is a nonsystemic, wide-spectrum organophosphate insecticide. These products are widely used against a variety of target pests on field crops, greenhouses and forests, as well as numerous indoor environments, from dog kennels to food processing plants, schools and hospitals.

Malathion is used to control aphids, mites, scale, flies, leafhoppers, leafminer, thrips, loopers, pear psylla, mealybugs, spittlebugs, corn earworms, chinchbugs, grasshoppers, armyworms, bollweevils, bollworms, lice, ticks, ants, spiders, and mosquitoes. It is applied to alfalfa, clover, pasture and range grasses, nonagricultural land, cereal crops, cotton, safflower, soybeans, sugar beets, corn, beans, blueberries, stored grain, and inside homes. Estimated use of malathion ranges from 10 to 15 million pounds of active ingredients used annually.

Health Effects

The U.S. Environmental Protection Agency (EPA) has classified malathion as a toxicity class III pesticide, bearing the signal word "Caution." Despite the fact that malathion is one of the less acutely toxic synthetic pesticides, having a rat oral LD50 of 1522 to 1945 milligrams per kilogram of body weight (mg/kg), numerous human poisonings have been reported. It is slightly toxic via the oral route and dermal route. Malathion is rapidly and effectively absorbed by practically all routes including the gastrointestinal tract, skin, mucous membranes, and lungs. Animal studies indicate it is eliminated through urine and feces with a reported half-life of approximately 8 hours in rats and approximately 2 days in cows.

Symptoms of acute exposure to organophosphate or cholinesterase-inhibiting compounds may include the following:

numbness, tingling sensations, headache, dizziness, tremor, nausea, abdominal cramps, sweating, incoordination, blurred vision, difficulty breathing or respiratory depression, and slow heartbeat. Very high doses may result in unconsciousness, incontinence, and convulsions or fatality. The antidote is atropine.

All organophosphate pesticides act on the body in the same way and their effects are additive, which raises questions about multiple exposure to these chemicals through many uses in and around homes and food production.

Malathion is a nerve poison, which acts by inhibiting the enzyme acetylcholine esterase (AChE), and probably acts at other sites in the nervous system as well. Cases of long-lasting polyneuropathy (Petty, 1958 and Healy, 1959), and sensory damage (Harrell, et al., 1978) have been reported in humans, as well as behavioral changes. Corresponding indications of neurotoxicity are seen in animal studies.

The pesticide has been shown in animal testing and from use experience to affect not only the central nervous system, but the immune system, adrenal glands, liver and blood.

Malathion has shown to be mutagenic in humans and animals. It has also been associated with birth defects in domestic and laboratory animals.

In a 1976 U.S. Army study, it was found that the avoidance behavior of rats was significantly impaired at a dose which did not impair blood or brain AChE activity. In other words, malathion was found to cause behavioral changes at levels at which the standard hospital test for organophosphate poisoning would be negative.

EPA's Pesticide Incident Monitoring System reported 962 incidents from 1960-80, and in California, malathion was the third most common cause of pesticide

related illness from 1981 - 85, especially among applicators exposed during indoor application, usually due to inhalation of fumes. Other physiological changes reported include glycosuria and hyperglycemia (sugar in the urine or blood) and elevated levels of polymorpholeucocytes (white blood cells).

A memo written by a Health Statistician with the Health Effects Division in the Office of Prevention, Pesticides and Toxic Substances, EPA, stated that 5,222 unintentional residential exposures were reported to Poison Control Centers from 1993-1996.

A November 1999 report issued by the Center for Disease Control (CDC) recommended that Florida find alternatives to malathion to fight the Mediterranean fruit fly (Medfly). According to the CDC report, over 230 people had reported being sick after malathion was sprayed aerially during the previous Medfly Eradication Program. A Florida Public Health Department study of people exposed to malathion has linked the chemical to respiratory, gastrointestinal and central nervous system effects.

Cancer Controversy

There is ongoing controversy about whether malathion is a carcinogen. A 1992 study published in *Cancer Research* linked the use of malathion by Iowa and Minnesota farmers to an increased risk of non-Hodgkin's lymphoma. Another study published in the *American Journal of Epidemiology* found similar increased risks in Nebraska farmers using malathion.

In a letter by Harold T. Smith, a U.S. Department of Agriculture official, to Dr. Robert Simon, executive director, Environmental and Toxicology International, a lab that tests for pesticide residues, regarding a draft Environmental Impact Statement (EIS) on fruit fly control procedures, Simon was told that the EIS was being held up due to EPA's ongoing review of

Beyond Pesticides

701 E Street, S.E., Suite 200 • Washington DC 20003
202-543-5450 (v) • 202-543-4791 (f)
info@beyondpesticides.org • www.beyondpesticides.org

malathion. Smith states, "Preliminary information received from EPA indicates that EPA is considering changing the registration status of malathion because of studies that suggest that it could be a low-level human carcinogen. Such a change would require review of protection and mitigation measures."

Just over a week before EPA released its preliminary risk assessment for malathion, the agency was poised to classify the organophosphate pesticide as a suspected carcinogen, according to Reuters newswire on May 10, 2000. Reuters's EPA source, speaking on condition of anonymity, revealed, "EPA scientists' risk assessment finds that malathion is a suspected carcinogen. The risks are fairly small, but acceptable compared to other substances." However, when the risk assessment was published in the May 11, 2000 *Federal Register* (Vol. 65, No. 92), EPA claimed there was "insufficient evidence" to assess the insecticide's cancer causing potential.

Over the past year, internal memos from an EPA senior toxicologist question EPA's reversal on its preliminary decision not to regulate the widely used insecticide malathion as a cancer causing agent. EPA reversed the cancer findings of independent scientists and adopted the conclusions of an industry panel, called the Pathology Working Group (PWG). The EPA toxicologist, in a memo to the Chairman of the Cancer Assessment Review Committee (CARC), concluded, "[U]nder EPA's guidelines in evaluating the tumorigenic response, the PWG report should be discounted, and the original diagnoses retained."

This is not the first time that malathion has been associated with a "cover-up" controversy. In 1978, the National Cancer Institute (NCI) funded five studies, which they concluded that malathion did not appear to be a carcinogen in experimental animals. Six years later, Dr. M. Adrian Gross, senior science advisor, Benefits and Use Division, Office of Pesticide Programs, EPA, reviewed the NCI studies and concluded that malathion does, in fact, behave as a carcinogen and presents an unacceptably high risk. His assessment of the data showed that malathion increased the rate of tumors in experimental animals, especially in the thyroid glands of

rats and mice. Gross further concluded that, "It has been established that the upper limit on the cancer risk attributable to currently established tolerances for malathion is as high as 1 to 4 per 1,000 population ... an unacceptably high limit."

Environmental Effects

Malathion is a non-specific poison, is known to be highly toxic to bees, and available data suggests that it is highly toxic to many aquatic non-target species, such as aquatic stages of amphibians, and aquatic invertebrates, and moderately toxic to birds. Malathion is toxic to many species of beneficial insects. It has a wide range of toxicities in fish, extending from very highly toxic in the walleye to highly toxic in brown trout and the cutthroat trout, and moderately toxic in fathead minnows.

Malathion has been determined to be the cause of the deaths of thousands of Staten Island fish by the New York Department of Environmental Conservation (DEC), as reported by the Queens Tribune. Tests conducted in two open water locations in Staten Island revealed that malathion is responsible for the deaths of close to 2,000 fish, said DEC spokesman Brian O'Conner. Preliminary tests found malathion levels in Willowbrook Lake water to be 5.8 part per billion (ppm) and 0.3 ppm in dead fish, says the Tribune.

Environmental Fate

Malathion is quite water-soluble (145 ppm at 25° C.), and preliminary data show that it is highly mobile in loam soils and thus a probable threat to groundwater. Malathion is of low persistence in soil with reported field half-lives of one to 25 days. Degradation in soil is rapid and related to the degree of soil binding. Breakdown occurs by a combination of biological degradation and nonbiological reaction with water. If released to the atmosphere, malathion will break down rapidly in sunlight, with a reported half-life in air of about 1.5 days. In river water, the half-life is less than one week, whereas malathion remained stable in distilled water for 3 weeks.

Toxic Contaminants

The Northwest Coalition for Alternatives

to Pesticides (NCAP) states that the toxicity of malathion is compounded by its metabolites and contaminants. Malaoxon, a metabolite produced by the oxidation of malathion in mammals, insects, plants, and in sunlight is a primary source of malathion's toxicity and is 40 times more acutely toxic than malathion. NCAP also states that over eleven chemical contaminants and analogues created in the production process have been found in technical malathion. These chemicals and the use of other pesticides can act synergistically with malathion to increase its toxicity.

Malathion Up for Review

Malathion is currently being evaluated by EPA to determine if current uses will remain on the market. As part of the ongoing implementation on the Food Quality Protection Act (FQPA), EPA has released the preliminary risk assessment for malathion and is requesting public comments, up to July 10, 2000.

The EPA's preliminary risk assessment for malathion can be viewed at : www.epa.gov/pesticides/op/malathion.htm.

The Federal Register announcement on malathion can be found at: www.epa.gov/fedrgstr/EPA-PEST/2000/May/Day-11/p11870.htm.

It is especially important for EPA to hear from people who have been poisoned by organophosphate pesticides. Please send written comments and/or fill out Beyond Pesticides/NCAMP's Pesticide Incident Record form and send it to EPA. Note on your comments: Control Number OPP-34223.

By mail, send your comments to: Public Information and Records Integrity Branch Information Resources and Services Division (7502C), Office of Pesticide Programs/EPA, 401 M Street, SW, Washington, DC 20460. By email, send your comments to: opp-docket@epa.gov. Note the above control number on the subject line. Electronic submissions will be accepted in WordPerfect 6.1/8.0 or ASCII file.

Revised July 2000

Beyond Pesticides

701 E Street, S.E., Suite 200 • Washington DC 20003
202-543-5450 (v) • 202-543-4791 (f)
info@beyondpesticides.org • www.beyondpesticides.org

Malathion *chemicalWATCH* Factsheet Bibliography

- Blondell, J. (1999). Memorandum to Susan Hummel Re: Review of Poison Control Center Data for Residential Exposures to Organophosphate Pesticides, 1993-1996. February 11th. Office of Prevention, Pesticides and Toxic Substances. Washington, DC.
- Brenner, L. (1992). "Malathion." *Journal of Pesticide Reform* 12(4).
- Cantor, K. et al. (1992). "Pesticides and other risk factors for non-Hodgkin's lymphomas among men in Iowa and Minnesota." *Cancer Research* 52:2447-2455.
- Deminti, B. (2000). Letter to William Burnam, Cancer Assessment Review Committee Chairman Re: evaluation of the carcinogenic potential of malathion. April 27th. Office of Pesticide Programs. Washington, DC.
- "EPA officials ignore the risk pesticide poses," *The Washington Post*, June 9, 1984.
- Extension Toxicology Network. (1996). "Malathion Pesticide Incident Profile." Oregon State University. <http://ace.orst.edu/info/extoxnet/>.
- Gosselin, R.E. (1984). *Clinical Toxicology of Commercial Products*. Williams & Wilkins, Baltimore, MD.
- Gross, A. (1984). Memorandum to Kevin Keaney Re: Carcinogenicity of malathion. April 24th. Office of Pesticide Programs. Washington, DC.
- Harell, et al. (1978). "Bilateral sudden deafness following combined insecticide poisoning." *Laryngoscope* 88:1348.
- Healy, J.K. (1959). "Ascending paralysis following malathion intoxication: A case report." *Med. J. Aust.* 1:765-767.
- Kurtz, P.J. (1976). "Behavioral and biochemical effects of malathion." Study No. 51-051-73/76. U.S. Army Environmental Hygiene Agency. Aberdeen Proving Ground, MD.
- Namba, T., et al. (1970). "Malathion poisoning: A fatal case with cardiac manifestations." *Arch. Environmental Health* 21:533-541.
- Petty, C. (1958). "Organic phosphate insecticide poisoning." *American Journal of Medicine* 24:467-470.
- "Respected researcher was fired over unpublished claims: Disputed study links malathion, cancer," *The Los Angeles Times*, September 17, 1981.
- Reuber, M. (1979). "Carcinogenicity of malathion." National Cancer Institute/Frederick Cancer Research Center. Frederick, MD.
- Thomson, W.T. (1984). *Agricultural Chemicals: Insecticides*. Thomson Publications, Fresno, CA.
- U.S. EPA. (2000). Evaluation of the Carcinogenic Potential of Malathion, attachments 1 through 29. Cancer Assessment Review Committee, Health Effects Division, Office of Pesticide Programs. Washington, DC.
- U.S. EPA. (1988). Guidance for the reregistration of pesticide products containing malathion as the active ingredient. Office of Pesticide Programs. Washington, DC.
- U.S. EPA. (1992). "Pesticide Industry Sales and Usage: 1990 and 1991 market estimates." Office of Pesticides and Toxic Substances. Washington, DC.
- U.S. EPA. (1980). "Summary of reported pesticide incidents involving malathion." Pesticide Incident Monitoring System (PIMS). Office of Pesticide Programs. Washington, DC.
- Weisenburger, D. et al. (1988). "A case-control study of non-Hodgkin's lymphoma and agricultural factors in eastern Nebraska." *American Journal of Epidemiology* 128:901.

Beyond Pesticides

701 E Street, S.E., Suite 200 • Washington DC 20003
202-543-5450 (v) • 202-543-4791 (f)
info@beyondpesticides.org • www.beyondpesticides.org